

Vertex reconstruction in scintillator

Haiqiong Zhang (from JUNO in China)

Yan Zhang (from JUNO in China)

Chiara Lastoria (from ProtoDUNE in Madrid, Spain)

Ryan Felkai (from NEXT in Valencia, Spain)

Carmen Romo Luque (from NEXT in Valencia, Spain)

2018.05.31 Mainz

Outline

- ▶ **Introduction of project**
- ▶ **Monte Carlo Simulation**
- ▶ **Vertex reconstruction**



Introduction

Solar neutrino detector

- ▶ Sphere $r = 5$ m
- ▶ Liquid scintillator
- ▶ PMTs



reconstruction algorithm for the position of the event vertex from the information extracted from the position and arrival time pattern



Monte Carlo simulation

Photon class

- ▶ Each photon is defined with a position of emission, a direction of propagation and a time of emission.

Event class

- ▶ Assuming they are point-like, an event is defined by its energy and its true position.
- ▶ Attributes of this class are the collection of photons and the collection of hits. Each hit consists of 3 spatial coordinates and the time of arrival.

Monte Carlo simulation

Physics class - Generation

- ▶ Takes event as an input, the first method is used to generate the photons.
- ▶ The number of photons is defined by the energy of the event and the light yield of the scintillator (600 photons per MeV).
- ▶ They are generated isotropically from the given position of the event.
- ▶ To simulate the characteristic of the light emission we generate at a time following an exponential decay (time constant 3 ns). Due to the deadline we set this emission time at 0 for all photons simplifying the problem drastically.

Monte Carlo simulation

Physics class - Propagation

- ▶ Takes also an event as an input.
- ▶ For each generated photon in the photon collection we compute the position of the hit on a sphere of 5m diameter.
- ▶ We make the reasonable assumptions that we have perfect transparency, infinite granularity, 100% photodetection probability and no light scattering.

Monte Carlo simulation


Physics class - Propagation

- ▶ Takes also an event as an input.
- ▶ For each generated photon in the photon collection we compute the position of the hit on a sphere of 5m diameter.
- ▶ We make the reasonable assumptions that we have perfect transparency, infinite granularity, 100% photodetection probability and no light scattering.



Monte Carlo simulation

Physics class - Propagation


- ▶ Takes also an event as an input.
 - ▶ For each generated photon in the photon collection we compute the position of the hit on a sphere of 5m diameter.
 - ▶ We make the reasonable assumptions that we have perfect transparency, infinite granularity, 100% photodetection probability and no light scattering.
- 
- ▶ The registered hit position and time of arrival are then registered and passed to our vertex reconstruction program₈

Generated events


For all the photons collected:

- Cartesian coordinates
- Spherical coordinates
- Arrival time on the sphere surface


Three situations



a) $P(0,0,0)$



b) $P(1,0,3)$




c) $P(0,4.75,0)$


Hit maps and arrival time

Hit maps

➤ $\phi:\sin\Theta$







photons arrival time
in the (ϕ, θ) plane







Generated events

a) $P(0,0,0)$







Generated events

b) $P(1,0,3)$



Generated events

c) $P(0, 4.75, 0)$



Vertex reconstruction

Algorithm principle

- When particles like e^+ , e^- or γ deposit energy in liquid scintillator, define the residual time:


$$t_{i,res} = t_i - tof - t_0$$

where:

t_i : first hit time of each PMT

t_{tof} : the time of flight for scintillator photon

t_0 : real time of an event, but in this project, it should be 0.



Vertex reconstruction

Algorithm principle

➤ For a certain scanned point \vec{R}_0 , calculate $t_{i,res}$

$$t_{i,res} = t_i - \frac{n * L(\vec{R}_0 - \vec{R}_i)}{c}$$

where n: refractive index of liquid scintillator;


c: light speed in vacuum;

\vec{R}_i : hit location in the detector, the total number is 600.

and define the parameter :

$$F(\vec{R}_0)_{\min} = \sum_{i=600} t_{i,res} * t_{i,res} \longrightarrow \vec{R}_0$$

\vec{R}_0 is the event vertex reconstructed by us.



Vertex reconstruction

Algorithm details

- create the cartesian coordinates

$$\theta \in (0, \pi)$$

$$\varphi \in (0, 2\pi)$$


$$r \in (0, 5)$$



$$x = r \sin\theta \cos\varphi$$

$$y = r \sin\theta \sin\varphi$$

$$z = r \cos\theta$$




- set the step for the loop of θ , φ , r , and get $\overrightarrow{R_0}$

$$\theta: \pi / 200$$

$$\varphi: \pi / 200$$

$$r: 5/200$$



Vertex reconstruction

Algorithm Results

For only one event:

reconstruction vertex

true vertex

difference

```
x=-1.55113e-15 y=4.2434e-16 z=-5
x=-1.5576e-15 y=3.99924e-16 z=-5
x=-1.56369e-15 y=3.75409e-16 z=-5
x=-1.56939e-15 y=3.50801e-16 z=-5
x=-1.57471e-15 y=3.26107e-16 z=-5
x=-1.57964e-15 y=3.01332e-16 z=-5
x=-1.58418e-15 y=2.76483e-16 z=-5
x=-1.58832e-15 y=2.51566e-16 z=-5
x=-1.59208e-15 y=2.26586e-16 z=-5
x=-1.59544e-15 y=2.01551e-16 z=-5
x=-1.59841e-15 y=1.76466e-16 z=-5
x=-1.60099e-15 y=1.51338e-16 z=-5
x=-1.60317e-15 y=1.26172e-16 z=-5
x=-1.60495e-15 y=1.00975e-16 z=-5
x=-1.60634e-15 y=7.5753e-17 z=-5
x=-1.60733e-15 y=5.05124e-17 z=-5
x=-1.60792e-15 y=2.52593e-17 z=-5
x=-1.60812e-15 y=-1.03442e-30 z=-5
X=0
Y=0
Z=0
root [1]
```

$(0,0,0)$

$(0,0,0)$

0m

```
x=-1.56939e-15 y=3.50801e-16 z=-5
x=-1.57471e-15 y=3.26107e-16 z=-5
x=-1.57964e-15 y=3.01332e-16 z=-5
x=-1.58418e-15 y=2.76483e-16 z=-5
x=-1.58832e-15 y=2.51566e-16 z=-5
x=-1.59208e-15 y=2.26586e-16 z=-5
x=-1.59544e-15 y=2.01551e-16 z=-5
x=-1.59841e-15 y=1.76466e-16 z=-5
x=-1.60099e-15 y=1.51338e-16 z=-5
x=-1.60317e-15 y=1.26172e-16 z=-5
x=-1.60495e-15 y=1.00975e-16 z=-5
x=-1.60634e-15 y=7.5753e-17 z=-5
x=-1.60733e-15 y=5.05124e-17 z=-5
x=-1.60792e-15 y=2.52593e-17 z=-5
x=-1.60812e-15 y=-1.03442e-30 z=-5
X=1.02034
Y=6.56333e-16
Z=2.98017
root [1]
```

$(1.02, 6.5E-16, 2.98)$

$(1,0,3)$

0.0284m

```
x=-1.54427e-15 y=4.48652e-16 z=-5
x=-1.55113e-15 y=4.2434e-16 z=-5
x=-1.5576e-15 y=3.99924e-16 z=-5
x=-1.56369e-15 y=3.75409e-16 z=-5
x=-1.56939e-15 y=3.50801e-16 z=-5
x=-1.57471e-15 y=3.26107e-16 z=-5
x=-1.57964e-15 y=3.01332e-16 z=-5
x=-1.58418e-15 y=2.76483e-16 z=-5
x=-1.58832e-15 y=2.51566e-16 z=-5
x=-1.59208e-15 y=2.26586e-16 z=-5
x=-1.59544e-15 y=2.01551e-16 z=-5
x=-1.59841e-15 y=1.76466e-16 z=-5
x=-1.60099e-15 y=1.51338e-16 z=-5
x=-1.60317e-15 y=1.26172e-16 z=-5
x=-1.60495e-15 y=1.00975e-16 z=-5
x=-1.60634e-15 y=7.5753e-17 z=-5
x=-1.60733e-15 y=5.05124e-17 z=-5
x=-1.60792e-15 y=2.52593e-17 z=-5
x=-1.60812e-15 y=-1.03442e-30 z=-5
X=-7.63858e-16
Y=-7.63858e-16
Z=-7.63858e-16
root [2]
```

$(-7.64E-16, 4.75, -7.64E-16)$

$(0, 4.75, 0)$

1.08×10^{-15}