The muon g - 2 and $\alpha(M_Z^2)$: the hadronic part

Fred Jegerlehner, DESY Zeuthen/Humboldt University Berlin

The Evaluation of the Leading Hadronic Contribution to the Muon Anomalous Magnetic Moment, Mainz Institute for Theoretical Physics, JG University Mainz, February 19-23, 2018



New updated and expanded edition Jegerlehner F., The Anomalous Magnetic Moment of the Muon. Springer Tracts in Modern Physics, Vol 274 (2017). Springer, Cham (693 pages on one number to 8 digits) Outline of Talk:

Introduction

alphi g - 2 Status: Theory vs experiment – do we see New Physics?

Hadronic Vacuum Polarization (HVP) – Data & Status

*****Evaluation of $\alpha(M_Z^2)$

 $\Delta \alpha^{had}$ Adler function controlled: Euclidean split trick

Issues in standard data based time-like approach

A problem in DR for HVP and a first direct measurement of $\Pi'_{\gamma}(s)$

Effective field theory: the Resonance Lagrangian Approach

HVP from lattice QCD

Alternative method: measure space-like $\alpha_{\text{QED,eff}}(t) \rightarrow a_{\mu}^{\text{had}}$

Conclusion

1. Introduction

Non-perturbative hadronic effects in electroweak precision observables, main effect via effective fine-structure "constant" $\alpha(E)$ (charge screening by vacuum polarization) Of particular interest:

 $\alpha(M_Z)$ and $a_\mu \equiv (g-2)_\mu/2 \Leftrightarrow \alpha(m_\mu)$

- electroweak effects (leptons etc.) calculable in perturbation theory
- ***** strong interaction effects (hadrons/quarks etc.) perturbation theory fails \implies Dispersion integrals over e^+e^- -data

encoded in

F. Jegerlehner

$$R_{\gamma}(s) \equiv \frac{\sigma(e^+e^- \to \gamma^* \to \text{hadrons})}{\sigma(e^+e^- \to \gamma^* \to \mu^+\mu^-)}$$

Errors of data \implies theoretical uncertainties !!! The art of getting precise results from non-precision measurements ! The challenge for precision experiments on $\sigma(e^+e^- \rightarrow hadrons)$ $\sigma_{hadronic}$ via scan or radiative return: e^+



NSK, KLOE, BABAR, BESIII...



Muon g - 2 to go

5 Numbers to establish the "g-2 Test"

(that is, 5 that have relevant uncertainties to keep watch on)



2. Muon g - 2 Status: Theory vs experiment

Given the CODATA/PDG recommended value of α the theory confronts experiment as follows: see Marc Knecht's Talk

Contribution	Value $\times 10^{10}$	Error×10 ¹⁰	Reference
QED incl. 4-loops + 5-loops	11 658 471.886	0.003	Aoyama et al 12,Laporta 17
Hadronic LO vacuum polarization	689.46	3.25	
Hadronic light-by-light	10.34	2.88	
Hadronic HO vacuum polarization	-8.70	0.06	
Weak to 2-loops	15.36	0.11	Gnendiger et al 13
Theory	11 659 178.3	3.5	-
Experiment	11 659 209.1	6.3	BNL 04
The Exp. 4.3 standard deviations	-30.6	7.2	-

Standard model theory and experiment comparison [in units 10^{-10}]. What represents the 4 σ deviation: \Box new physics? \Box a statistical fluctuation? \Box underestimating uncertainties (experimental, theoretical)? \diamond do experiments measure what theoreticians calculate?

- a "New Physics" interpretation of the persisting 3 to 4 σ requires relatively strongly coupled states in the range below about 250 GeV.
- Search bounds from LEP, Tevatron and specifically from the LHC already have ruled out a variety of Beyond the Standard Model (BSM) scenarios, so much hat standard motivations of SUSY/GUT extensions seem to fall in disgrace.
- There is no doubt that performing doable improvements on both the theory and the experimental side allows one to substantially sharpen (or diminish) the apparent gap between theory and experiment.
- * Or is it unaccounted for real photon radiation effects? Do experiments measure what theory calculates?
- At the present/future level of precision a_{μ} depends on all physics incorporated in the SM: electromagnetic, weak, and strong interaction effects and beyond that all possible new physics we are hunting for.

here we are and hope to go:



Past and future g - 2 experiments testing various contributions. New Physics $\stackrel{?}{=}$ deviation $(a_{\mu}^{exp} - a_{\mu}^{the})/a_{\mu}^{exp}$. Limiting theory precision: hadronic vacuum polarization (HVP) and hadronic light-by-light (HLbL)

*** digging deeper and deeper ***

same status for the electron:



Status and sensitivity of the a_e experiments testing various contributions. The error is dominated by the uncertainty of $\alpha(\text{Rb11})$ from atomic interferometry. No "New Physics" $\stackrel{?}{=}$ deviation $(a_e^{\exp} - a_e^{\text{the}})/a_{\mu}^{\exp}$. The blue band illustrates the improvement by the Harvard experiment. Note the very different sensitivities to non-QED contributions in comparison with a_{μ} .

*** still is and remains a QED test mainly ***

Experiment now:

$a_{\mu}^{\exp} = (11\,659\,209.1\pm 5.4\pm 3.3[6.3]) \times 10^{-10}$ BNL updated

To come – :

New muon g - 2 experiments at Fermilab and J-PARC: improve error by factor 4

 \Rightarrow new muon g - 2 experiment: $\Delta a_{\mu} = a_{\mu}^{exp} - a_{\mu}^{the} = 8 \sigma$ theory as today

Reduction of hadronic VP uncertainty by factor 2, same HLbL $\Rightarrow \Delta a_{\mu} = 12 \sigma$

That's what we hope to achieve!

Key problem: limited accuracy of HVP! How to safely reduce and crosscheck?

New lattice QCD now starts to have impact

Ongoing improvement on R measurements

New alternative methods : the workshops main theme here

F. Jegerlehner

R-data Evaluation of a_{μ}^{had}



F. Jegerlehner

muonLOHCws@MITP, JGU Mainz, February 19-23, 2018

Issues in standard data based time-like approach

How to combine a pretty large number of data-sets to a truly reliable *R*-function: true uncertainty? How much just taken from pQCD? Choosing/selecting data-sets? Bare vs physical cross sections, how reliable is VP subtraction?

2 Radiative corrections specifically for ISR method, sQED issues etc. The ISR method requires one order in α more precise RC calculation relative to SCAN method, at least full 2–loop Bhabha and/or $e^+e^- \rightarrow \mu^+\mu^-$ as well as ISR–FSR interference in $\pi^+\pi^-$ channel. What about RC to other more complicated channels (see e.g. F.J.&Karol Kołodziej 2017)? see Hernryk Czyż's Talk What about disentangling 30 channels and recombining them in the 1 to 2 GeV region (quantum interference, missing parts, double counting issues)?

What precisely do we need in the DR? The 1PI "blob", which is not a measurable quantity. Need undressing from QED effects, photon VP subtraction, FSR modeling, $\rho^0 - \gamma$ mixing? Do we do this at sufficient precision?

On-convergence of Dyson series for OZI suppressed narrow resonances

In addition:



Recent BES-III vs BaBar and KLOE: dominant $\pi\pi$ channel still could be better

Recent results:

• $\pi^+\pi^-$ from BES-III, CMD-3 and CLEOc • $\pi^+\pi^-\pi^0$ from Belle • K^+K^- from CMD-3 and SND • $\omega\pi^0 \to \pi^0\pi^0\gamma$ from SND • $K_S K^{\pm}\pi^0\pi^{\mp}, K_S K^{\pm}\pi^{\mp}\eta, \pi^+\pi^-\pi^0\pi^0, K_S K_L\pi^0, K_S K_L\eta, K_S K_L\pi^0\pi^0$ from BaBar

see Simon Eidelman's Talk

Energy range	a_{μ}^{had} [%](error) × 10 ¹⁰	rel. err.
$ ho, \omega~(E < 1~{ m GeV})$, 540.98 [78.6](2.80)	0.5 %
1 GeV < E < 2 GeV	96.49 [14.0](2.54)	2.6 %
2 GeV < $E < \infty$ incl pQCD	51.09 [7.4](1.10)	2.2 %
total	688.65 [100.0](3.94)	0.6 %



abs

50

41

100

Still an issue in HVP

region 1.2 to 2 GeV data; test-ground exclusive vs inclusive R measurements (more than 30 channels!) VEPP-2000 CMD-3, SND (NSK) scan, BaBar, BES III radiative return! still contributes 50% of uncertainty



illustrating progress by BaBar and NSK exclusive channel data vs new inclusive data by KEDR. Why point at 1.84 GeV so high?

□ A problem in DR for HVP?

Full photon propagator Dyson resummation of 1PI part (blue blob)

1

$$\begin{split} \gamma & \gamma \\ & \gamma \\$$

Including external e.m. coupling

$$e^2 D'_{\gamma}(q^2) = \frac{-i}{q^2} \frac{e^2}{1 + \Pi'_{\gamma}(q^2)}$$

Effective charge

F. Jegerlehner

$$\frac{e^2}{1+\Pi_{\gamma}'(s)} = \frac{e^2}{1-\Delta\alpha(s)} = e^2(s)$$

Usually, $\Delta \alpha(s)$ is a correction i.e $\Delta \alpha(s) \ll 1$ and the Dyson series converges well.

Exceptions: narrow OZI suppressed resonances (below $q\bar{q}$ -thresholds)



 Γ_{ee} not much smaller than Γ_{QCD} (i.e strong decays): $J/\psi, \psi_2, \Upsilon_1, \Upsilon_2, \Upsilon_3$

Note: imaginary parts from narrow resonances, $\lim \Pi'(s) = \frac{\alpha}{3} R(s) = \frac{3}{\alpha} \frac{\Gamma_{ee}}{\Gamma}$ at peak, are sharp spikes and are obtained correctly only by appropriately high resolution scans. For example,

$$|1 - \Pi'(s)|^2 - (\alpha/\alpha(s))^2 = (\mathsf{Im} \ \Pi'(s))^2$$

F. Jegerlehner

at $\sqrt{s} = M_R$ is given by

ρ	1.23	$\times 10^{-3}$
ω	2.76	$\times 10^{-3}$
ϕ	1.56	$\times 10^{-2}$
J/ψ	594.81	
ψ_2	9.58	
ψ_3	2.66	$\times 10^{-4}$
Υ_1	104.26	
Υ_2	30.51	
Υ_3	55.58	

• What is measured in an experiment is the full propagator, corresponding to $\frac{1}{1-x}; x \text{ irreducible part}$ Object required in the DR: $R_{\gamma}(s) \equiv \sigma^{(0)}(e^+e^- \rightarrow \gamma^* \rightarrow \text{hadrons}) / \frac{4\pi\alpha^2}{3s} R^{\text{undressed}} = R^{\text{observed}} * |(1-x)|^2$

VP subtraction is iterative procedure: does not converge!

The time-like vs space-like effective charge



Note that the smooth space-like effective charge agrees rather well with the non-resonant "background" above the Φ (kind of duality)

No proof that this cannot produce non-negligible shifts!

Time-like VP-subtraction cannot be implemented locally near OZI suppressed resonances: J/ψ , ψ' and Υ_1 , Υ_2 , Υ_3

Present leading uncertainty: hard to improve by direct R(s) measurements



Comparison with other Results. Note: results depend on which value is taken for HLbL. JS11 and BDDJ13 includes $116(39) \times 10^{-11}$ [JN], DHea09, DHMZ10, HLMNT11 and BDDJ12 use $105(26) \times 10^{-11}$ [PdRV].

HVP from lattice QCD

The need for ab initio calculation of a_{μ}^{had} is well motivated:

- the problems to determine non-perturbative contributions to the muon g 2 from experimental data at sufficient precision persists and is not easy to improve,
- a model—independent extension of CHPT to the relevant energies ranges up to 2 GeV is missing while the new experiments E989 FNAL and E34 J-PARC

require an improvement of the hadronic uncertainties by a factor of four.

The hope is that LQCD can deliver estimates of accuracy

$$\delta a_{\mu}^{\mathrm{HVP}}/a_{\mu}^{\mathrm{HVP}} < 0.5\%$$
, $\delta a_{\mu}^{\mathrm{HLbL}}/a_{\mu}^{\mathrm{HLbL}} \lesssim 10\%$

in the coming years.

see Marina Marinkovic's Talk

F. Jegerlehner

Primary object for HVP in LQCD: e.m. current correlator in configuration space

$$\langle J_{\mu}(\vec{x},t) J_{\nu}(\vec{0},0) \rangle$$
, $J_{\mu} = \frac{2}{3}\bar{u}\gamma_{\mu}u - \frac{1}{3}\bar{d}\gamma_{\mu}d - \frac{1}{3}\bar{s}\gamma_{\mu}s + \cdots$

In principle, a Fourier transform

$$\Pi_{\mu\nu}(Q) = \int \mathrm{d}^4 x \mathrm{e}^{\mathrm{i}\,Qx} \left\langle J_{\mu}(x) \, J_{\nu}(0) \right\rangle = \left(Q_{\mu} Q_{\nu} - \delta_{\mu\nu} \, Q^2 \right) \, \Pi(Q^2)$$

yields the vacuum polarization function $\Pi(Q^2)$ needed to calculate

$$a_{\mu}^{\rm HVP} = 4\alpha^2 \int_0^\infty dQ^2 f(Q^2) \left\{ \Pi(Q^2) - \Pi(0) \right\}$$

The integration kernel in this representation is

$$f(Q^2) = w(Q^2/m_{\mu}^2)/Q^2$$
; $w(r) = \frac{16}{r^2(1+\sqrt{1+4/r})^4\sqrt{1+4/r}}$.



The integrand Q^2 . Ranges between $Q_i = 0.00, 0.15, 0.30, 0.45$ and 1.0 GeV and their percent contribution to a_{μ}^{had} and the "LQCD sample"

□ LQCD lattice in finite box: momenta are quantized Q_{min} = 2π/L where L is the lattice box length. Q_{min} → 0 ⇔ L → ∞ infinite volume limit
 □ Q_{min} = 2π/L with m_πaL ≥ 4 for m_π ~ 200 MeV, such that Q_{min} ~ 314 MeV
 □ about 44% of the low x contribution to a^{had}_u is not covered by data yet



• lattice data: $Q^2 > (2\pi/L)^2$

- \Rightarrow extrapolate to $Q^2 = 0$ via Padé's
- ♦ Note need $\Pi(0)$!
- ✤ required accuracy: needed LQCD data down to $Q^2_{min} \approx 0.1 \text{ GeV}^2$

New: RBC/UKQCD 18 use lattice between 0.1 and 4 GeV² and *R*-data for IR and UV tails \Rightarrow most precise evaluation so far: $a_{\mu}^{\text{HVP-LO}} = (692.5 \pm 2.7) \times 10^{-10}$ $[a_{\mu}^{\text{lat}} = (232.1 \pm 1.5) \times 10^{-10} \stackrel{\circ}{=} 33.5\%$; $a_{\mu}^{\text{dat}} = (460.4 \pm 2.2) \times 10^{-10}$]



Summary of recent LQCD results for the leading order a_u^{HVP} , in units 10^{-10} . Labels: \blacksquare marks $u, d, s, c, \blacktriangle u, d, s$ and $I_{u,d}$ contributions. Individual flavor contributions from light (u, d)amount to about 90%, strange about 8% and charm about 2%. Budapest, Marseille, Wuppertal, Brookhaven, Zeuthen, Mainz, Edinburgh, ... The gray vertical band represents my evaluation. The wheat band represents the HVP required such that theory matches the experimental BNL result. The very precise RBC/UKQCD point is obtained by supplementing lattice results by R-data.

3. $\alpha(M_Z^2)$ in precision physics (precision physics limitations)

Uncertainties of hadronic contributions to effective α are a problem for electroweak precision physics:

 $\begin{array}{ll} \alpha \ , \ G_{\mu}, M_{Z} \ \text{most precise input parameters} \\ 50\% \ \text{non-perturbative} \end{array} \Rightarrow \begin{array}{l} \text{precision predictions} \\ \sin^{2}\Theta_{f}, v_{f}, a_{f}, M_{W}, \Gamma_{Z}, \Gamma_{W}, \cdots \end{array}$

 $\alpha(M_Z), G_\mu, M_Z$ best effective input parameters for VB physics (Z,W) etc.

$$\frac{\delta \alpha}{\sigma_{\mu}} \sim 3.6 \times 10^{-9}$$

$$\frac{\delta G_{\mu}}{G_{\mu}} \sim 8.6 \times 10^{-6}$$

$$\frac{\delta M_Z}{M_Z} \sim 2.4 \times 10^{-5}$$

$$\frac{\delta \alpha(M_Z)}{\alpha(M_Z)} \sim 0.9 \div 1.6 \times 10^{-4}$$
 (present : lost 10⁵ in precision!)
$$\frac{\delta \alpha(M_Z)}{\alpha(M_Z)} \sim 5.3 \times 10^{-5}$$
 (ILC requirement)

LEP/SLD:
$$\sin^2 \Theta_{\text{eff}} = (1 - g_{Vl}/g_{Al})/4 = 0.23148 \pm 0.00017$$

 $\delta \Delta \alpha(M_Z) = 0.00020 \qquad \Rightarrow \qquad \delta \sin^2 \Theta_{\text{eff}} = 0.00007$

affects Higgs mass bounds, precision tests and new physics searches!!!

For pQCD contributions very crucial: precise QCD parameters α_s , m_c , m_b , $m_t \Rightarrow$ Lattice-QCD

muonLOHCws@MITP, JGU Mainz, February 19-23, 2018



parameter relationships between very precisely measurable quantities precision tests, possible sign of new physics non-perturbative $\Delta \alpha_{had}^{(5)}(M_Z^2)$ is limiting precision predictions

Note: 30 SD disagreement between SM prediction and experiment when subleading corrections are dropped!

SM extrapolation up to Planck scale?

After Higgs discovery: Higgs vacuum stability issue! \Rightarrow Need very precise SM parameters: g', g, g_s, y_t, λ



The SM dimensionless couplings in the $\overline{\text{MS}}$ scheme as a function of the renormalization scale for $M_H = 124 - 127$ GeV.

perturbation expansion works up to the Planck scale!

no Landau pole or other singularities, Higgs potential likely remains stable! $\Box U(1)_Y$ screening (IR free), $SU(2)_L$, $SU(3)_c$ antiscreening (UV free): g_1, g_2, g_3

as expected (standard wisdom)

Top Yukawa y_t and Higgs λ : screening if standalone (IR free, like QED)

as part of SM, transmutation from IR free to UV free

As SM couplings are as they are: QCD dominance in top Yukawa RG requires $g_3 > \frac{3}{4}y_t$, top Yukawa dominance in Higgs RG requires $\lambda < \frac{3(\sqrt{5}-1)}{2}y_t^2$ in the gaugeless $(g_1, g_2 = 0)$ limit.

In the focus:

 \Box does Higgs self-coupling stay positive $\lambda > 0$ up to Λ_{Pl} ?

The key question/problem concerns the size of the top Yukawa coupling y_t decides about stability of our world! — [$\lambda = 0$ would be essential singularity!]

Will be decided by:
more precise input parameters

better established EW matching conditions

I R-data Evaluation of $\alpha(M_Z^2)$

Non-perturbative hadronic contributions $\Delta \alpha_{had}^{(5)}(s) = -(\Pi_{\gamma}'(s) - \Pi_{\gamma}'(0))$ can be evaluated in terms of $\sigma(e^+e^- \rightarrow \text{hadrons})$ data via dispersion integral: $\Delta \alpha_{\text{had}}^{(5)}(s) = -\frac{\alpha s}{3\pi} \left(\iint_{\gamma}^{E_{\text{cut}}^2} ds' \frac{R_{\gamma}^{\text{data}}(s')}{s'(s'-s)} \right)$ $+ \underbrace{\mathcal{F}}_{E_{\text{cut}}^{2}}^{\infty} ds' \frac{R_{\gamma}^{\text{pQCD}}(s')}{s'(s'-s)} \right)$ Compilation: FJ 15 Theory = pQCD: Gorishny et al. 91, Chetyrkin et al. 97...09 $J/\psi_{1S} \psi_{2S}$ e'e' -> hadrons DQCD $R_{\gamma}(s) \equiv \frac{\sigma^{(0)}(e^+e^- \to \gamma^* \to \text{hadrons})}{\frac{4\pi\alpha^2}{3s}}$ where 3.1 GeV 2.0 GeV œ3 $1.0 \ {\rm GeV}$ 5.2 GeV ϕ • M3N ρ, ω ∗ BĒ $\circ \gamma \gamma 2$ BESI $0.0 \ {\rm GeV}, \infty$ 0 $9.5~{\rm GeV}$ 1.0 2.5 3.5 1.5 2.0 3.0 4.0 4.5 5.0 E (GeV) p-QCD 13.GeV had T15 T25 35 45 e'e' -> hadrons $\sim \sigma_{\rm tot}^{\rm had}(q^2)$ $\Pi_{\gamma}^{'\,\mathrm{had}}(q^2)$ hadronic vacuum polarization с CB • JADE -S- pQCD × MD-1 TASSO DHHM 13 11 $\alpha(s) = \frac{\alpha}{1 - \Delta \alpha(s)}$; $\Delta \alpha(s) = \Delta \alpha_{\text{lep}}(s) + \Delta \alpha_{\text{had}}^{(5)}(s) + \Delta \alpha_{\text{top}}(s)$ E (GeV)

Present situation: (after KLOE, BaBar and first BESIII results)

$\Delta \alpha_{\rm hadrons}^{(5)}(M_Z^2)$	=	0.027738 ± 0.000158	
		0.027523 ± 0.000119	Adler
$\alpha^{-1}(M_Z^2)$	=	128.919 ± 0.022	
		128.958 ± 0.016	Adler

$\Delta \alpha_{\rm had}(M_Z^2)$ results from ranges:

for $M_Z = 91.1876$ GeV in units 10^{-4} . 2017 update in terms of e^+e^- -data and pQCD. 43% data, 57% perturbative QCD. pQCD is used between 5.2 GeV and 9.5 GeV and above 11.5 GeV.

final state	range (GeV)	$\Delta \alpha_{\rm had}^{(5)} imes 10^4$ (stat) (syst) [tot]	rel	abs
ρ	(0.28, 1.05)	33.91 (0.05) (0.18)[0.19]	0.6%	1.4%
ω	(0.42, 0.81)	3.10 (0.04) (0.08)[0.09]	3.0%	0.3%
ϕ	(1.00, 1.04)	4.76 (0.07) (0.11)[0.13]	2.7%	0.7%
J/ψ		12.38 (0.60) (0.67)[0.90]	7.2%	32.1%
Ŷ		1.30 (0.05) (0.07)[0.09]	6.9%	0.3%
had	(1.05, 2.00)	16.53 (0.06) (0.83)[0.83]	5.0%	27.4%
had	(2.00, 3.20)	15.34 (0.08) (0.61)[0.62]	4.0%	15.2%
had	(3.10, 3.60)	4.98 (0.03) (0.09)[0.10]	1.9%	0.4%
had	(5.20, 5.20)	16.84 (0.12) (0.21)[0.25]	0.0%	2.4%
pQCD	(5.20, 9.46)	33.84 (0.12) (0.25)[0.03]	0.1%	0.0%
had	(9.46,11.50)	11.12 (0.07) (0.69)[0.69]	6.2%	19.2%
pQCD	(11.50,∞)	123.29 (0.00) (0.05)[0.05]	0.0%	0.1%
data	(0.28,11.50)	120.25 (0.63) (1.45)[1.58]	1.0%	0.0%
total		277.38 (0.63) (1.45)[1.58]	0.6%	100.0%

Correlation between different contributions to $a_{\mu}^{\rm had}$ and $\Delta \alpha^{\rm had\,(5)}$



Contributions from e^+e^- data ranges and form pQCD to a_{μ}^{had} and $\Delta \alpha^{had(5)}$.

4. Reducing uncertainties via the Euclidean split trick: Adler function controlled pQCD

experiment side: new more precise measurements of *R*(*s*)
 future direct measurements Patrick Janot, Luca Trentadue et al
 theory side: $\alpha_{\rm em}(M_7^2)$ by the "Adler function controlled" approach

$$\alpha(M_Z^2) = \alpha^{\text{data}}(-s_0) + \left[\alpha(-M_Z^2) - \alpha(-s_0)\right]^{\text{pQCD}} + \left[\alpha(M_Z^2) - \alpha(-M_Z^2)\right]^{\text{pQCD}}$$

where the space-like $-s_0$ is chosen such that pQCD is well under control for $-s < -s_0$. The monitor to control the applicability of pQCD is the Adler function

$$D(Q^{2} = -s) = \frac{3\pi}{\alpha} s \frac{d}{ds} \Delta \alpha_{\text{had}}(s) = -(12\pi^{2}) s \frac{d\Pi'_{\gamma}(s)}{ds} = Q^{2} \int_{4m^{2}}^{\infty} \frac{R(s)}{(s+Q^{2})^{2}}$$

which also is determined by R(s) and can be evaluated in terms of experimental e^+e^- -data. Perturbative QCD tail: $D(Q^2) \rightarrow N_c \sum_f Q_f^2 (1 + O(\alpha_s))$ as $Q^2 \rightarrow \infty$.

S. Eidelman, F. J., A. Kataev, O. Veretin, Phys. Lett. B 454 (1999) 369

$\Delta \alpha^{had}$ Adler function controlled

✓ use old idea: Adler function: Monitor for comparing theory and data

$$D(-s) \doteq \frac{3\pi}{\alpha} s \frac{d}{ds} \Delta \alpha_{\text{had}}(s) = -\left(12\pi^2\right) s \frac{d\Pi_{\gamma}'(s)}{ds}$$
$$\Rightarrow \quad D(Q^2) = Q^2 \left(\int_{4m_{\pi}^2}^{E_{\text{cut}}^2} \mathrm{d}s \frac{R(s)^{\text{data}}}{(s+Q^2)^2} + \int_{E_{\text{cut}}^2}^{\infty} \frac{R^{\text{pQCD}}(s)}{(s+Q^2)^2} \,\mathrm{d}s\right).$$

$\mathbf{pQCD} \leftrightarrow R(s)$	$\mathbf{pQCD} \leftrightarrow D(Q^2)$		
very difficult to obtain	smooth simple function		
in theory	in <u>Euclidean</u> region		

Conclusion:

★time-like approach: pQCD works well in "perturbative windows" 3.00 - 3.73 GeV, 5.00 - 10.52 GeV and 11.50 - ∞ Kühn,Steinhauser ★space-like approach: pQCD works well for $\sqrt{Q^2 = -q^2} > 2.0$ GeV (see plot) "Experimental" Adler–function versus theory (pQCD + NP)

Error includes statistical + systematic here (in contrast to most *R*-plots showing statistical errors only)!



(Eidelman, F. J., Kataev, Veretin 98, FJ 08/17 updates) theory based on results by Chetyrkin, Kühn et al.

 \Rightarrow pQCD works well controlled to predict $D(Q^2)$ down to $s_0 = (2.0 \text{ GeV})^2$; use this to calculate

$$\Delta \alpha_{\rm had}(-Q^2) \sim \frac{\alpha}{3\pi} \int \mathrm{d}Q'^2 \frac{D(Q'^2)}{Q'^2}$$

$$\Delta \alpha_{\rm had}^{(5)}(-M_Z^2) = \left[\Delta \alpha_{\rm had}^{(5)}(-M_Z^2) - \Delta \alpha_{\rm had}^{(5)}(-s_0) \right]^{\rm pQCD} + \Delta \alpha_{\rm had}^{(5)}(-s_0)^{\rm data}$$

and obtain, for $s_0 = (2.0 \text{ GeV})^2$:

$$\Delta \alpha_{had}^{(5)}(-s_0)^{data} = 0.006409 \pm 0.000063$$
$$\Delta \alpha_{had}^{(5)}(-M_Z^2) = 0.027483 \pm 0.000118$$
$$\Delta \alpha_{had}^{(5)}(M_Z^2) = 0.027523 \pm 0.000119$$

♦ shift +0.000008 from the 5-loop contribution
♦ error ±0.000100 added in quadrature form perturbative part
QCD parameters: • $\alpha_s(M_Z) = 0.1189(20)$,

• $m_c(m_c) = 1.286(13) [M_c = 1.666(17)]$ GeV, • $m_b(m_c) = 4.164(25) [M_b = 4.800(29)]$ GeV

based on a complete 3–loop massive QCD analysis Kühn et al 2007

F. J., Nucl. Phys. Proc. Suppl. 181-182 (2008) 135

(FJ 98/17)

$\Delta \alpha_{\rm had}(-M_0^2)$ results from ranges:

for $M_0 = 2$ GeV in units 10^{-4} . 2015 update in terms of e^+e^- -data and pQCD. 94% data, 6% perturbative QCD. pQCD is used between 5.2 GeV and 9.5 GeV and above 11.5 GeV.

final state	range (GeV)	$\Delta \alpha_{\rm had}^{(5)}(-M_0^2) \times 10^4$ (stat) (syst) [tot]	rel	abs
ρ	(0.28, 1.05)	29.78 (0.04) (0.16)[0.16]	0.5%	6.6%
ω	(0.42, 0.81)	2.69 (0.03) (0.07)[0.08]	3.0%	1.6%
ϕ	(1.00, 1.04)	3.78 (0.05) (0.09)[0.10]	2.7%	2.6%
J/ψ		3.21 (0.15) (0.15)[0.21]	6.7%	11.4%
Y		0.05 (0.00) (0.00)[0.00]	6.8%	0.0%
had	(1.05, 2.00)	10.36 (0.04) (0.49)[0.49]	4.8%	61.2%
had	(2.00, 3.20)	6.06 (0.03) (0.25)[0.25]	4.2%	16.1%
had	(3.10, 3.60)	1.31 (0.01) (0.02)[0.03]	1.9%	0.2%
had	(5.20, 5.20)	2.90 (0.02) (0.02)[0.03]	0.0%	0.2%
pQCD	(5.20, 9.46)	2.66 (0.02) (0.02)[0.00]	0.1%	0.0%
had	(9.46,11.50)	0.39 (0.00) (0.02)[0.02]	5.7%	0.1%
pQCD	(11.50,∞)	0.90 (0.00) (0.00)[0.00]	0.0%	0.0%
data	(0.28,11.50)	60.53 (0.18) (0.61)[0.63]	1.0%	0.0%
total		64.09 (0.18) (0.61)[0.63]	1.0%	100.0%

Of $\Delta \alpha_{had}^{(5)}(M_Z^2)$ 22% data, 78% pQCD!



Contributions from e^+e^- data ranges and form pQCD to $\Delta \alpha_{had}^{(5)}(-M_0^2)$ vs. $\Delta \alpha_{had}^{(5)}(M_Z^2)$.

	$[86\%,\!13\%]$			
	$[52\%,\!47\%]$			
	$[57\%,\!42\%]$			
	$[18\%,\!81\%]$			
	$[84\%,\!15\%]$			
	$[84\%,\!15\%]$			
	$[56\%,\!43\%]$			
	$[16\%,\!83\%]$			
	$[84\%,\!15\%]$			
	$[29\%,\!70\%]$			
	$[20\%,\!79\%]$			
	$[20\%,\!79\%]$			
	$[56\%,\!43\%]$			
	$[54\%,\!45\%]$			
	$[38\%,\!41\%]$			
	$[26\%,\!73\%]$			
	[50%, 49%]			
	$[29\%,\!70\%]$			
	$[45\%,\!54\%]$			
	[21%,77%]			
20	40	60	80	1(

 $[\Delta \alpha_{\rm had}^{\rm data} / \Delta \alpha_{\rm had}^{\rm tot}, \Delta \alpha_{\rm had}^{\rm pQCD} / \Delta \alpha_{\rm had}^{\rm tot}]$ in %

How much pQCD?

Jegerlehner 1985 Lynn et al. 1985 Burkhardt et al. 1989 Martin, Zeppenfeld 1994 Swartz 1995 Eidelman, Jegerlehner 1995 Burkhardt, Pietrzyk 1995 Adel, Yndurain 1995 Alemany, Davier, Höcker 1997 Kühn, Steinhauser 1998 Davier, Höcker 1998 **Erler 1998** Burkhardt, Pietrzyk 2001 Hagiwara et al 2004 Jegerlehner 2006 direct Jegerlehner 2006 Adler Hagiwara et al. 2011 Davier et al. 2011 Jegerlehner 2016 direct Jegerlehner 2016 Adler data-driven
theory-driven
fifty-fifty
low energy weighted data

Note: the Adler function monitored Euclidean data vs pQCD split approach is only moderately more pQCD-driven, than the time-like approach adopted by Davier et al. and others.

Alternative method: measure space-like $\alpha_{\text{QED,eff}}(t)$

Newly proposed recently: [arXiv:1504.02228,1609.08987]

"A new approach to evaluate the leading hadronic corrections to the muon g-2" Carloni Calame, Passera, Trentadue, Venanzoni 2015; Abbiendi et al. 2016

□ space-like
$$\Delta \alpha_{had}(-Q^2) = 1 - \frac{\alpha}{\alpha(-Q^2)} - \Delta \alpha^{lep}(-Q^2)$$
 determines a_{μ}^{had} via
$$a_{\mu}^{had} = \frac{\alpha}{\pi} \int_{0}^{1} dx (1-x) \Delta \alpha_{had} \left(-Q^2(x)\right)$$

where $Q^2(x) \equiv \frac{x^2}{1-x}m_{\mu}^2$ is the space–like square momentum–transfer. Also in the Euclidean region the integrand is highly peaked, now around half of the ρ meson mass scale.

F. Jegerlehner



The integrand of a_{μ}^{had} integral as functions of *x* and *Q*. Strongly peaked at about 330 MeV. Ranges between $Q_i = 0.00, 0.15, 0.30, 0.45$ and 1.0 GeV and their percent contribution to a_{μ}^{had} .

The measuring directly low energy running $\alpha_{\text{QED}}(s)$ in space-like region via

- very different paradigm: no VP subtraction issue!
- no exclusive channel collection
- even 1% level measurement can provide important independent information

Bhabha scattering





VP dressed tree level Bhabha scattering in QED

has two tree level diagrams the *t*- and the *s*-channel. With the positive c.m. energy square $s = (p^+ + p^-)^2$ and the negative momentum transfer square

$$t = (p_{-} - p'_{-})^{2} = -\frac{1}{2} (s - 4m_{e}^{2}) (1 - \cos \theta),$$

 θ the e^- scattering angle, there are two very different scales involved

The VP dressed lowest order cross-section is

$$\frac{\mathrm{d}\sigma}{\mathrm{d}\cos\Theta} = \frac{s}{48\pi} \sum_{ik} |A_{ik}|^2$$

where A_{ik} tree level helicity amplitudes, i, k = L, R left– and right–handed electrons.

Dressed transition amplitudes: $(m_e \approx 0)$

$$|A_{\text{LL,RR}}|^2 = \frac{3}{8} (1 + \cos \theta)^2 \left| \frac{e^2(s)}{s} + \frac{e^2(t)}{t} \right|^2$$
$$|A_{\text{LR,RL}}|^2 = \frac{3}{8} (1 - \cos \theta)^2 \left| \frac{e^2(s)}{s} + \frac{e^2(t)}{t} \right|^2$$

Preferably one uses small angle Bhabha scattering (small |t|) as a normalizing process which is dominated by the *t*-channel ~ 1/t, however, detecting electrons and positrons along the beam axis often has its technical limitations.

F. Jegerlehner

Care also is needed concerning the ISR corrections because cuts for the Bhabha process $(e^+e^- \rightarrow e^+e^-)$ typically are different from the ones applied to $e^+e^- \rightarrow$ hadrons. Usually, experiments have included corresponding uncertainties in their systematic errors, if they not have explicitly accounted for all appropriate radiative corrections.



Get a_{μ}^{had} from $\mu^-e^- \rightarrow \mu^-e^-$ process

G. Abbiendi et al., arXiv:1609.08987

Massimo Passera's Talk

$$\frac{\mathrm{d}\sigma_{\mu^- e^- \to \mu^- e^-}^{\mathrm{unpol.}}}{\mathrm{d}t} = 4\pi \,\alpha(t)^2 \,\frac{1}{\lambda(s, m_e^2, m_\mu^2)} \,\left\{ \frac{\left(s - m_\mu^2 - m_e^2\right)^2}{t^2} + \frac{s}{t} + \frac{1}{2} \right\}$$

• The primary goal of [arXiv:1504.02228,1609.08987]: determining a_{μ}^{had} in an alternative way

•
$$\Pi'_{\gamma}(Q^2) - \Pi'_{\gamma}(0) = -\Delta \alpha_{had}(-Q^2) = \frac{\alpha}{\alpha(-Q^2)} + \Delta \alpha^{lep}(-Q^2) - 1$$

directly checks lattice QCD data

My proposal here: determine very accurately

 $\Delta \alpha_{\rm had} \left(-Q^2\right)$ at $Q \approx 2.5 \; {\rm GeV}$

by this method (one single number!) as the non-perturbative part of $\Delta \alpha_{had} \left(M_Z^2 \right)$ as in "Adler function" approach.

Conclusions

- Muon g 2 theory uncertainty remains the key issue.
- Are presently estimated (essentially agreed) evaluations in terms of *R*-data reliable?
- Lattice QCD estimates very close to be competitive, tending to larger central values?
- Novel hybrid method lattice + R-data optimized method looks very promising D. Bernecker, Harvey B. Meyer 2011, RBC/UKQCD Blum et al. 2018
- In any case on paper $e^-\mu^+ \rightarrow e^-\mu^+$ looks to be the ideal process to perform an unambiguous measurement of $\alpha(-Q^2)$, which determines the LO HVP to a_μ
- Radiative corrections much easier than for the time-like hadronic channels
- Key problem: how to control in a fixed target experiment the precision at the 1% level? What range is accessible at what precision?

Likely to be supplemented by lattice QCD results and/or using time-like data

This experiment G. Abbiendi et al. is absolutely important also as it allows for for direct crosschecks with lattice QCD results and is has completely different systematics. Even a 5% crosscheck would be very helpful to scrutinize the HVP issue, and last bu not least whether the observed deviation is a real BSM effect.

Thanks you for your attention!