



Power Expansions on the Light-cone @ MITP, September 21, 2022

Factorization, Resummation and Endpoint Singularities in Higgs to Diphoton Decay

Ze Long Liu

u^b

b
UNIVERSITÄT
BERN

Motivation


- Form factor of meson

Brodsky, Lepage, PRD 22 (1980) 2157

$$F(Q^2) \sim \frac{1}{Q} \left[\int dx \int dy T(x, y, \mu) \phi_a(x, \mu) \phi_b(y, \mu) + \mathcal{O}(1/Q) \right]$$

at leading power in $1/Q$

$$\phi_\pi \propto x(1-x)f_\pi \quad T(x, y) = \frac{8\pi\alpha_s(Q)}{9xy}$$



Phenomenology important !

convolution is well-defined at leading power

Power suppressed hard exchange tend to give contributions diverging as $\int \frac{dx}{x}$

Bauer, Pirjol, Stewart, '02, Rothstein, '03

overviews by Martin and Gherardo

- Soft-Collinear Effective Theory (SCET) admit a consistent power expansion
 - Soft quark field is involved at subleading-power Lagrangian
 - Endpoint-divergent convolutions arise from soft quark


Moult, Stewart, Vita, Zhu et al., Beneke, Garny, Szafron, Wang et al., 2018-2020

Subleading Power

- A simple scale separation is violated by endpoint divergences at subleading power
- Failure of $\overline{\text{MS}}$ subtraction scheme
 - $h \rightarrow \gamma\gamma$ mediated by b-quark loop is a simple example:
no non-perturbative effect
- Evaluating the scattering amplitudes in the high energy limit is a fundamental problem of QFT
- Resummation of large logarithms is necessary for a reliable theoretical prediction

Leading vs. Subleading

Leading Power



- Factorization is local


$$H(Q^2, \mu) J_n(M, \mu, \nu/Q) J_{\bar{n}}(M, \mu, \nu/Q) S(M, \mu, \nu/M)$$

$$\sim 1 + \frac{\alpha}{4\pi} (-L^2 + 4L + c_0) + \mathcal{O}(\alpha^2) \quad L = \ln \frac{-Q^2}{M^2}$$

- No convolution, no endpoint

- rapidity divergence

Subleading Power



$$\lambda \sim \frac{m_b}{M_h}$$

$$L = \ln \frac{-M_h^2}{m_b^2}$$

- Factorization is non-local

$$H_1 O_1 + H_2 \otimes O_2 + H_3 \otimes J \otimes J \otimes S$$

$$\sim y_b \boxed{m_b} \left\{ \left(\frac{L^2}{2} - 2 \right) + \frac{C_F \alpha_s}{4\pi} \left[-\frac{L^4}{12} - L^3 + \dots \right] \right\} + \mathcal{O}(\lambda^2)$$

- Endpoint-divergent convolutions

- rapidity divergence

Rapidity divergences at LO

$$\int d^d \ell \frac{1}{(\ell^2 - m^2)(n \cdot \ell + i\epsilon)(-\bar{n} \cdot \ell + i\epsilon)}$$

$$\sim \int_{m_b^2}^{\infty} \frac{d(\ell^+ \ell^-)}{(\ell^+ \ell^-)} (\ell^+ \ell^- - m_b^2)^{-\epsilon} \int_0^{\infty} \frac{d\ell^+}{\ell^+}$$


↑
divergence at $\ell^+ \ell^- \rightarrow +\infty$
is dimensionally regulated

Additional regulator is needed !!!

analytical regulator: $\left(\frac{\nu}{\ell^+}\right)^\eta$ Becher, Bell, '11

rapidity regulator: $\left|\frac{\ell^+ - \ell^-}{\nu}\right|^\eta$ Chiu, Jain, Neill, Rothstein, '12

exponential regulator: $e^{-\tau(\ell^+ + \ell^-)}$ Li, Neill, Zhu, '16



soft mode is scaleless
rapidity div. cancel between col. & anti-col.

symmetric with $\ell^+ \leftrightarrow \ell^-$
non-trivial soft mode

Rapidity divergences at LO

We prefer a regulator:

- gives non-trivial soft function
- symmetric with $\ell^+ \leftrightarrow \ell^-$
- easy for higher order calculation

ZLL, Neubert, 1912.08818(JHEP)

$$\Theta(\ell^+ - \ell^-) \left(-\frac{M_h \ell^+}{\nu^2} \right)^\eta + \Theta(\ell^- - \ell^+) \left(-\frac{M_h \ell^-}{\nu^2} \right)^\eta$$

hard $\frac{1}{\epsilon^2} - \frac{L_h}{\epsilon} + \frac{L_h^2}{2} - 2 - \frac{\pi^2}{12}$ $L_h = \ln \frac{-M_h^2}{\mu^2}$

col.+ anti-col. $\left(\frac{2}{\eta} + 2 \ln \frac{-M_h^2}{\nu^2} \right) \left(\frac{1}{\epsilon} - L_m \right)$ $L_m = \ln \frac{m_b^2}{\mu^2}$



soft $- \left(\frac{2}{\eta} + \ln \frac{-M_h^2}{\nu^2} + \ln \frac{m_b^2}{\nu^2} \right) \left(\frac{1}{\epsilon} - L_m \right) - \frac{1}{\epsilon^2} + \frac{L_m}{\epsilon} - \frac{L_m^2}{2} + \frac{\pi^2}{12}$

$H + C + \bar{C} + S = \frac{L^2}{2} - 2$ $L = \ln \frac{-M_h^2}{m_b^2}$

Factorization Theorem at Subleading Power and Endpoint Divergences

ZLL, M. Neubert, 1912.08818 (JHEP)

Factorization at Bare Level




- perfect scale separation with $z \sim \mathcal{O}(1)$ and $\ell^+, \ell^- \sim \mathcal{O}(m_b)$
- factorization violated at $z \ll 1, \ell^-/\ell^+ \gg 1$ $\ell^+, \ell^- \sim M_h$

Factorization at Bare Level


$$\begin{aligned}\mathcal{M}(h \rightarrow \gamma\gamma) = & H_1 O_1 + 4 \int_0^1 dz \left(-\frac{z M_h^2}{\nu^2} \right)^\eta H_2(z) O_2(z) \\ & + 2 H_3 \int_0^\infty \frac{d\ell_-}{\ell_-} \int_0^{\ell_-} \frac{d\ell_+}{\ell_+} \left(-\frac{M_h \ell_-}{\nu^2} \right)^\eta J(M_h \ell_+) J(-M_h \ell_-) S(\ell_+ \ell_-, m_b^2) \\ & + (\ell_+ \leftrightarrow \ell_-)\end{aligned}$$

- each ingredient is well defined with dimensional regulator
- regulator is friendly for higher order calculations
- all the divergences cancel, but highly non-trivial !!!
- divergent convolution gives $\frac{1}{\eta}$ and $\frac{1}{\epsilon}$ poles

Matrix Elements




$$\langle \gamma\gamma | O_1 | h \rangle = \frac{m_b}{e_b^2} \langle \gamma\gamma | h(0) \mathcal{A}_{n_1}^{\perp\mu}(0) \mathcal{A}_{n_2,\mu}^{\perp}(0) | h \rangle$$



$$\langle \gamma\gamma | O_2(z) | h \rangle = \langle \gamma\gamma | h(0) \left[\bar{\chi}_{n_1}(0) \gamma_{\perp}^{\mu} \frac{\not{n}_1}{2} \delta(z M_h + i \bar{n}_1 \cdot \partial) \chi_{n_1}(0) \right] \mathcal{A}_{n_2,\mu}^{\perp}(0) | h \rangle$$

$$\lim_{z \rightarrow 0} O_2(z) = \frac{1}{\epsilon} - L_m + \frac{C_F \alpha_s}{4\pi} \left(\frac{m_b^2}{\mu^2} \right)^{-\epsilon} \text{ non-vanish at } z=0, \text{ factorization violated}$$

$$\times \left\{ \frac{1}{\epsilon^2} (\ln z + 3) + \frac{1}{\epsilon} \left(\frac{\ln^2 z}{2} - \frac{1}{2} - \frac{\pi^2}{6} \right) + \frac{\ln^3 z}{6} + \left(\frac{\pi^2}{6} - \frac{1}{2} \right) \ln z + \frac{\pi^2}{6} - 2\zeta_3 + 3 \right\}$$




$$\langle \gamma\gamma | O_3 | h \rangle = \langle \gamma\gamma | T \left\{ h(0) \bar{\chi}_{n_1}(0) \chi_{n_2}(0), i \int d^D x \mathcal{L}_{q \xi_{n_1}}^{(1/2)}(x), i \int d^D y \mathcal{L}_{\xi_{n_2} q}^{(1/2)}(y) \right\} | h \rangle$$

$$\mathcal{L}_{q \xi_{n_1}}^{(1/2)}(x) = \bar{q}_s(x_-) [\mathcal{A}_{n_1}^{\perp}(x) + \mathcal{G}_{n_1}^{\perp}(x)] \chi_{n_1}(x)$$

coupling of soft quark to hard-collinear fields

Factorization of O_3 at Operator Level

$$\mathcal{L}_{q \xi_{n_1}}^{(1/2)}(x) = \bar{q}_s(x_-) [\mathcal{A}_{n_1}^\perp(x) + \mathcal{G}_{n_1}^\perp(x)] \mathcal{X}_{n_1}(x)$$



$$\langle \gamma\gamma | O_3 | h \rangle = \langle \gamma\gamma | T \left\{ h(0) \bar{\chi}_{n_1}(0) \chi_{n_2}(0), i \int d^D x \mathcal{L}_{q \xi_{n_1}}^{(1/2)}(x), i \int d^D y \mathcal{L}_{\xi_{n_2} q}^{(1/2)}(y) \right\} | h \rangle$$

hard-col. quark field $\chi_{n_i}(x) = \frac{\not{p}_i \not{\ell}_i}{4} W_{n_i}^\dagger(x) b(x)$

decouple with
soft mode $\chi_{n_1}(x) \rightarrow S_{n_1}(x_-) \chi_{n_1}(x),$
 $\mathcal{G}_{n_1}^{\perp\mu}(x) \rightarrow S_{n_1}(x_-) \mathcal{G}_{n_1}^{\perp\mu}(x) S_{n_1}^\dagger(x_-)$

collinear & soft Wilson line

$$W_{n_i}(x) = P \exp \left[ig_s \int_{-\infty}^0 ds \bar{n}_i \cdot G_{n_i}(x + s\bar{n}_i) \right]$$

$$S_{n_i}(x) = P \exp \left[ig_s \int_{-\infty}^0 dt n_i \cdot G_s(x + tn_i) \right]$$


hard-collinear sector

$$O_3 = h(0) \int d^D x \int d^D y T \left\{ [(\mathcal{A}_{n_1}^\perp(x) + \mathcal{G}_{n_1}^\perp(x)) \chi_{n_1}(x)]^{\alpha i} \bar{\chi}_{n_1}^{\beta j}(0) \right\} \cdot T \left\{ \chi_{n_2}^{\beta k}(0) [\bar{\chi}_{n_2}(y) (\mathcal{A}_{n_2}^\perp(y) + \mathcal{G}_{n_2}^\perp(y))]^{\gamma l} \right\}$$

hard-anti-collinear sector

$$\times T \left\{ [S_{n_2}^\dagger(y_+) q_s(y_+)]^{\gamma l} [\bar{q}_s(x_-) S_{n_1}(x_-)]^{\alpha i} [S_{n_1}^\dagger(0) S_{n_2}(0)]^{jk} \right\} \text{ soft sector}$$

Radiative Jet Function



$$\langle \gamma(k_1) | T \left[(\mathcal{A}_{n_1}^\perp(x) + \mathcal{G}_{n_1}^\perp(x)) \mathcal{X}_{n_1}(x) \right] \bar{\chi}_{n_1}(r) | 0 \rangle \sim \int \frac{d^D p}{(2\pi)^D} \frac{i \bar{n}_1 \cdot p}{p^2 + i0} J(p^2, (p - k_1)^2) e^{-ip \cdot (x-r) + ik_1 \cdot x}$$

light-like


Renormalization: $J(p^2, \mu) = \int_0^\infty dx Z_J(p^2, xp^2; \mu) J^{(0)}(xp^2)$ reach endpoint region

First obtained in $B^- \rightarrow \gamma \ell^- \bar{\nu}$ Bosch, Hill, Lange, Neubert, '03

	Local	
$Z_J(p^2, xp^2; \mu) = \left[1 + \frac{C_F \alpha_s}{4\pi} \left(-\frac{2}{\epsilon^2} + \frac{2}{\epsilon} \ln \frac{-p^2}{\mu^2} \right) \right] \delta(1-x) +$	Non-local	$\frac{C_F \alpha_s}{2\pi \epsilon} \Gamma(1, x) + \mathcal{O}(\alpha_s^2)$

with $\Gamma(y, x) = \left[\frac{\theta(y-x)}{y(y-x)} + \frac{\theta(x-y)}{x(x-y)} \right]_+$ Lange-Neubert Kernel

Soft-Quark Soft Function



$$\frac{i}{2} \mathcal{S}(\ell_+ \ell_-) P_n = -\frac{(4\pi)^{1-\epsilon}}{N_c} e^{\epsilon \gamma_E} \int dr_1 e^{ir_1 \ell_-} \int dr_2 e^{-ir_2 \ell_+} \\ \times \mu^{2\epsilon} \langle 0 | T \text{Tr } \bar{P}_{\bar{n}} S_{\bar{n}}(0, r_1 \bar{n}) q_s(r_1 \bar{n}) \bar{q}_s(r_2 n) S_n(r_2 n, 0) P_n | 0 \rangle$$

$$S(\ell_+ \ell_-) = \frac{1}{2\pi i} [\mathcal{S}(\ell_+ \ell_- + i0) - \mathcal{S}(\ell_+ \ell_- - i0)] \quad \mathcal{S}(\ell_+ \ell_-) \equiv \int \frac{d^{D-2} \ell_\perp}{(2\pi)^{D-2}} \mathcal{S}_1(\ell)$$

$$S(\ell_+ \ell_-) \sim m_b [S_a(\ell_+ \ell_-) \theta(\ell_+ \ell_- - m_b^2) + S_b(\ell_+ \ell_-) \theta(m_b^2 - \ell_+ \ell_-)]$$

Reproduce NLO amplitude

$$\mathcal{M}(h \rightarrow \gamma\gamma) = H_1 O_1 \quad \dots \quad T_1$$

$$+ 4 \int_0^1 dz \left(-\frac{z M_h^2}{\nu^2} \right)^\eta H_2(z) O_2(z) \quad \dots \quad T_2$$

$$+ 2H_3 \int_0^\infty \frac{d\ell_-}{\ell_-} \int_0^{\ell_-} \frac{d\ell_+}{\ell_+} \left(-\frac{M_h \ell_-}{\nu^2} \right)^\eta J(M_h \ell_+) J(-M_h \ell_-) S(\ell_- \ell_+, m_b^2) \quad \dots \quad T_3$$

$$T_1 = \frac{1}{\epsilon^2} - \frac{L_h}{\epsilon} + \frac{L_h^2}{2} - 2 - \frac{\pi^2}{12} + \frac{C_F \alpha_s}{4\pi} \left(-\frac{1}{2\epsilon^4} + \dots \right),$$

$$T_2 = \left[\frac{2}{\eta} + 2 \ln \frac{-M_h^2 - i0}{\nu^2} \right] \left[\frac{1}{\epsilon} - L_m + \frac{C_F \alpha_s}{4\pi} \left(-\frac{3}{\epsilon^3} + \dots \right) \right] + \frac{C_F \alpha_s}{4\pi} \left(-\frac{2}{\epsilon^4} + \dots \right),$$

$$T_3 = - \left[\frac{2}{\eta} + \ln \frac{-M_h^2 - i0}{\nu^2} + \ln \frac{m_b^2}{\nu^2} \right] \left[\frac{1}{\epsilon} - L_m + \frac{C_F \alpha_s}{4\pi} \left(-\frac{3}{\epsilon^3} + \dots \right) \right] \\ - \frac{1}{\epsilon^2} + \frac{L_m}{\epsilon} - \frac{L_m^2}{2} + \frac{\pi^2}{12} + \frac{C_F \alpha_s}{4\pi} \left(\frac{5}{2\epsilon^4} + \dots \right)$$

NLO amplitude is perfectly reproduced up to constant terms

because of endpoint divergences, the poles in T_2 and T_3
are higher than the poles in $H_2(z), O_2(z), J(p^2), S(\ell^+ \ell^-)$

Endpoint Divergences

$\mathcal{O}(\alpha_s)$	$\lim_{z \rightarrow 0} H_2(z)$	$\lim_{z \rightarrow 0} O_2(z)$
LO	$\frac{1}{z}$	$\frac{1}{\epsilon} - L_m$
NLO	$\frac{c_0}{z} + \frac{c_1}{z^{1+\epsilon}}$	$a_0 + a_1 z^\epsilon$
NNLO	$\frac{c'_0}{z} + \frac{c'_1}{z^{1+\epsilon}} + \frac{c'_2}{z^{1+2\epsilon}}$	$a'_0 + a'_1 z^\epsilon + a'_2 z^{2\epsilon}$

Can we use plus expansion to resolve endpoint divergences?

$$z^{-1+\eta} = \frac{1}{\eta} \delta(z) + \left(\frac{1}{z} \right)_+ + \eta \left(\frac{\ln z}{z} \right)_+ + \dots$$

undefined convolution arises

$$\int_0^1 dz \left(\frac{1}{z} \right)_+ \ln z$$

difficult to renormalize endpoint divergences before performing convolution !!!

Refactorization

At the endpoint, the refactorization conditions read

Talk by Philipp

$$[\![H_2(z)]\!] \equiv \lim_{z \rightarrow 0} H_2(z) = H_3 \frac{J(z M_h^2)}{z}$$

$$[\![O_2(z)]\!] \equiv \lim_{z \rightarrow 0} \langle \gamma\gamma | O_2(z) | h \rangle = \int_0^\infty \frac{d\ell_+}{\ell_+} J(M_h \ell_+) S(\ell_+ \ell_-),$$

Then $[\![H_2(z)]\!] \otimes [\![O_2(z)]\!]$ has the same integrand with $H_3 O_3$,
but has different integral region


$$\int_0^1 dz [\![H_2(z)]\!] [\![O_2(z)]\!] = H_3 \int_0^{M_h} \frac{d\ell^-}{\ell^-} J(M_h \ell^-) \int_0^\infty \frac{d\ell^+}{\ell^+} J(M_h \ell^+) S(\ell^+ \ell^-)$$

useful to do subtraction

$$\int_0^1 dz H_2(z) O_2(z) \rightarrow \int_0^1 dz (H_2(z) O_2(z) - [\![H_2(z)]\!] [\![O_2(z)]\!]) + \int_0^1 dz [\![H_2(z)]\!] [\![O_2(z)]\!]$$

Refactorization at Operator Level

ZLL, Neubert, Mecaj, Wang, '20




$$H_3^{(0)} \langle b(k_1) \bar{b}(zk_1) \gamma(k_2) | T \left\{ h \bar{\chi}_{n_1} \chi_{n_2}(0), i \int d^D y \mathcal{L}_{\xi_{n_2} q}^{(1/2)}(y) \right\} | h \rangle$$

$$H_3^{(0)} \langle b(k_1) \bar{b}(zk_1) \gamma(k_2) | T \left\{ \bar{\chi}_{n_1}(0) S_{n_1}^\dagger(0) S_{n_2}(0) \chi_{n_2}(0) \right.$$

$$\times i \int d^D y \bar{\chi}_{n_2}(y) (\mathcal{A}_{n_2}^\perp(y) + \mathcal{G}_{n_2}^\perp(y)) S_{n_2}^\dagger(y_+) q_s(y_+) \left. \right\} | 0 \rangle$$

$$[\![H_2^{(0)}(z)]\!] = \frac{[\![\bar{H}_2^{(0)}(z)]\!]}{z} = -\frac{H_3^{(0)}}{z} J(z M_h^2)$$



$$O_2^{(0)}(t) = h(0) \bar{\chi}_{n_1}(0) \gamma_\perp^\mu \frac{\not{p}_1}{2} \chi_{n_1}(t \bar{n}_1) \mathcal{A}_{n_2, \mu}^\perp(0)$$

$$h(0) \bar{\chi}_{n_1}(0) S_{n_1}^\dagger(0) S_{n_2}(0) \gamma_\perp^\mu \frac{\not{p}_2}{2} S_{n_2}^\dagger(t n_2) q_s(t n_2) \mathcal{A}_{n_2, \mu}^\perp(0)$$


$$[\![\langle \gamma \gamma | O_2^{(0)}(z) | h \rangle]\!] = -\frac{g_\perp^{\mu\nu}}{2} \int_0^\infty \frac{d\ell_+}{\ell_+} J^{(0)}(-M_h \ell_+) S^{(0)}(z M_h \ell_+)$$

Result in two same integrands between $H_2 O_2$ and $H_3 O_3$

Cancellation of Rapidity Divergence

$$\int_0^a dz \frac{1}{z} + \int_b^{+\infty} dz \frac{1}{z} = \int_0^a dz \frac{1}{z^{1-\eta}} + \int_b^{+\infty} dz \frac{1}{z^{1-\eta}} = \frac{1}{\eta} (a^\eta - b^\eta) = \ln \frac{a}{b}$$

$$\int_0^a \frac{dz}{z} f(z) + \int_b^{+\infty} \frac{dz}{z} f(z) \rightarrow \int_b^a \frac{dz}{z} f(z)$$



$$H_3 \left(\int_{M_h}^{\infty} d\ell^- \int_0^{\infty} d\ell^+ + \int_0^{M_h} d\ell^- \int_0^{\infty} d\ell^+ \right) J(M_h \ell^-) J(M_h \ell^+) S(\ell^+ \ell^-) = 0$$

$\int_0^1 dz [H_2(z)] [O_2(z)]$

rapidity divergences cancel explicitly !!!


Rearrangement

$$S_\infty(\ell^+ \ell^-) = \lim_{\ell^\pm \rightarrow \infty} S(\ell^+ \ell^-, m_b^2)$$

$$S_\infty(\ell_+ \ell_-) \sim m_b \left(\frac{\ell_+ \ell_-}{\mu^2} \right)^{-\epsilon} \left[\frac{e^{\epsilon \gamma_E}}{\Gamma(1-\epsilon)} + \frac{C_F \alpha_s}{4\pi} \left(\frac{\ell_+ \ell_-}{\mu^2} \right)^{-\epsilon} \left(-\frac{2}{\epsilon^2} + \frac{6}{\epsilon} + 12 - \frac{\pi^2}{3} \right) \right]$$

$S_\infty(\ell^+ \ell^-)$ is not at soft scale, but at hard scale

$$\begin{aligned} \Delta H_1 O_1 &= -H_3 \lim_{\sigma \rightarrow -1} \int_{M_h}^{\infty} \frac{d\ell_-}{\ell_-} \int_{\sigma M_h}^{\infty} \frac{d\ell_+}{\ell_+} J(M_h \ell_-) J(-M_h \ell_+) S_\infty(\ell_+ \ell_-) \\ &\quad + (\ell_+ \leftrightarrow \ell_-) \end{aligned}$$



$$\begin{aligned} \mathcal{M}(h \rightarrow \gamma\gamma) &= (H_1 + \Delta H_1) O_1 \\ &\quad + 4 \lim_{\delta \rightarrow 0} \int_{\delta}^1 dz \left[H_2(z) O_2(z) - [\![H_2(z)]\!] [\![O_2(z)]\!] \right] \\ &\quad + H_3 \lim_{\sigma \rightarrow -1} \int_0^{M_h} \frac{d\ell_-}{\ell_-} \int_0^{\sigma M_h} \frac{d\ell_+}{\ell_+} J(M_h \ell_-) J(-M_h \ell_+) S(\ell_+ \ell_-) + (\ell_+ \leftrightarrow \ell_-) \end{aligned}$$

get rid of rapidity regulator !!!

no endpoint divergences any more !!!

Rearrangement

$$\begin{aligned} \mathcal{M}(h \rightarrow \gamma\gamma) &= (H_1 + \Delta H_1) O_1 && T_1 \\ &+ 4 \lim_{\delta \rightarrow 0} \int_\delta^1 dz \left[H_2(z) O_2(z) - [\![H_2(z)]\!] [\!O_2(z)\!]\right] && T_2 \\ &+ H_3 \lim_{\sigma \rightarrow -1} \int_0^{M_h} \frac{d\ell_-}{\ell_-} \int_0^{\sigma M_h} \frac{d\ell_+}{\ell_+} J(M_h \ell_-) J(-M_h \ell_+) S(\ell_+ \ell_-) && T_3 \end{aligned}$$

$$T_1 = -2 + \frac{C_F \alpha_s}{4\pi} \left[-\frac{\pi^2}{3\epsilon^2} + \frac{1}{\epsilon} \left(\frac{2\pi^2}{3} L_h - 10\zeta_3 \right) - \frac{2\pi^2}{3} L_h^2 + (12 + 20\zeta_3) L_h - 36 - \frac{7\pi^4}{30} \right],$$

$$T_2 = \frac{C_F \alpha_s}{4\pi} \left[\frac{\pi^2}{3\epsilon^2} + \frac{1}{\epsilon} \left(-\frac{2\pi^2}{3} L_h + 2\zeta_3 \right) + \frac{\pi^2}{3} (L_h^2 + 2L_h L_m - L_m^2) - 4\zeta_3 L_h + 8\zeta_3 + \frac{13\pi^4}{90} \right],$$

$$T_3 = \frac{L^2}{2} + \frac{C_F \alpha_s}{4\pi} \left[\frac{8\zeta_3}{\epsilon} - \frac{L^4}{12} - L^3 + \left(4 - \frac{\pi^2}{3} \right) L^2 - \left(8 - \frac{2\pi^2}{3} \right) L \right. \\ \left. + L_m (-3L^2 + 6L - 16\zeta_3) - 4\zeta_3 - \frac{\pi^4}{9} \right]$$

- NLO amplitude is exactly reproduced !!!
 - T_1 only depends on hard log
 - no $1/\epsilon^4$ poles any more
 - LL and NLL only appear in T_3
 - T_3 is “almost” RG-invariant

Resummation at LL

Leading Log only appears in the “almost” RG-invariant T_3

Sudakov exponent of the integrand:

$$H_3(\mu) J(M_h \ell_-, \mu) J(-M_h \ell_+, \mu) S(\ell_+ \ell_-, \mu) \sim m_b \theta(\ell_+ \ell_- - m_b^2) \exp \left[-\frac{C_F \alpha_s}{2\pi} \ln \frac{M_h}{\ell_-} \ln \frac{-M_h - i0}{\ell_+} + \dots \right]$$

Performing convolution with cutoff:

$$\begin{aligned} \mathcal{M}_b(h \rightarrow \gamma\gamma)|_{\text{LL}} &= \lim_{\sigma \rightarrow -1} \mathcal{M}_0 \int_0^{\sigma M_h} \frac{d\ell_+}{\ell_+} \int_0^{M_h} \frac{d\ell_-}{\ell_-} \theta(\ell_+ \ell_- - m_b^2) \exp \left[-\frac{C_F \alpha_s}{2\pi} \ln \frac{M_h}{\ell_-} \ln \frac{\sigma M_h}{\ell_+} \right] \\ &= \mathcal{M}_0 \frac{L^2}{2} {}_2F_2 \left(1, 1; \frac{3}{2}, 2; -\frac{C_F \alpha_s}{8\pi} L^2 \right) \quad \text{T. Liu, Penin, '17} \quad \text{J. Wang, '19} \end{aligned}$$

Mismatch in Cutoff Scheme

T_3 is “RG-invariant”:

$$T_3 = H_3 \int_0^\infty \frac{d\ell_-}{\ell_-} \int_0^\infty \frac{d\ell_+}{\ell_+} J(M_h \ell_-) J(-M_h \ell_+) S(\ell_+ \ell_-) \quad \text{bare}$$

$$= H_3(\mu) \int_0^\infty \frac{d\ell_-}{\ell_-} \int_0^\infty \frac{d\ell_+}{\ell_+} J(M_h \ell_-, \mu) J(-M_h \ell_+, \mu) S(\ell_+ \ell_-, \mu) \quad \text{renormalized}$$

This can be employed to determine Z-factor of soft function, Talk by Xiangpeng
but the convolution is not well defined!

When cutoff is involved:

$$T_3 = H_3 \int_0^{M_h} \frac{d\ell_-}{\ell_-} \int_0^{M_h} \frac{d\ell_+}{\ell_+} J(M_h \ell_-) J(-M_h \ell_+) S(\ell_+ \ell_-) \quad \text{bare}$$

$$\neq H_3(\mu) \int_0^{M_h} \frac{d\ell_-}{\ell_-} \int_0^{M_h} \frac{d\ell_+}{\ell_+} J(M_h \ell_-, \mu) J(-M_h \ell_+, \mu) S(\ell_+ \ell_-, \mu) \quad \text{renormalized}$$

$$\begin{aligned} \delta T_3 &\sim \int_0^{\sigma M_h} \frac{d\ell'_+}{\ell'_+} \left(\int_0^\infty d\ell_- \int_0^{M_h} d\ell'_- - \int_0^{M_h} d\ell_- \int_0^\infty d\ell'_- \right) \theta(\ell'_+ \ell'_- - m_b^2) \\ &\times (\ell'_+ \ell'_- - m_b^2)^{-\epsilon} \left[\frac{\theta(\ell'_- - \ell_-)}{\ell'_- (\ell'_- - \ell_-)} + \frac{\theta(\ell_- - \ell'_-)}{\ell_- (\ell_- - \ell'_-)} \right]_+ \neq 0 \end{aligned}$$

A Short Summary

- Factorization theorem is built at bare level
- NLO amplitude is reproduced exactly
- Non-trivial soft-quark soft function is derived
- Refactorization helps us to regulate the rapidity divergences
- Endpoint divergences are removed by rearrangement

How to deal with the mismatch between renormalization
and convolutions with cutoffs?

Renormalized Factorization Theorem at Subleading Power in SCET

ZLL, Neubert, 2003.03393 (JHEP)

ZLL, Mecaj, Neubert, Wang, Fleming, 2005.03013 (JHEP)

ZLL, M. Neubert, B. Mecaj, X. Wang

2009.04456 (PRD) & 2009.06779 (JHEP)

Renormalized Factorization Theorem

T1: Absorb the contributions from infinity-bin and mismatch

$$\mathcal{M}_b = H_1(\mu) \langle O_1(\mu) \rangle$$

T2: Divergences at the endpoints are subtracted

$$+ 2 \int_0^1 dz \left[H_2(z, \mu) \langle O_2(z, \mu) \rangle - \frac{\llbracket \bar{H}_2(z, \mu) \rrbracket}{z} \llbracket \langle O_2(z, \mu) \rangle \rrbracket - \frac{\llbracket \bar{H}_2(\bar{z}, \mu) \rrbracket}{\bar{z}} \llbracket \langle O_2(\bar{z}, \mu) \rangle \rrbracket \right]$$

$$+ g_{\perp}^{\mu\nu} H_3(\mu) \lim_{\sigma \rightarrow -1} \int_0^{M_h} \frac{d\ell_-}{\ell_-} \int_0^{\sigma M_h} \frac{d\ell_+}{\ell_+} J(M_h \ell_-, \mu) J(-M_h \ell_+, \mu) S(\ell_+ \ell_-, \mu) \Big|_{\text{leading power}}$$

T3: Cutoff is involve to regulated endpoint divergences

- Renormalization does NOT commute with the cutoff → **Mismatch**
- Cutoff arises from rearrangement – **Nature !**
- Both of the contributions from infinity-bin and mismatch only depend on hard scale – **Proved to all order in α_s**
- Renomalization of O_1 and O_2 is mixing

Renormalization

$$H_1(\mu) \sim H_1^{(0)} Z_{11}^{-1} + H_2^{(0)}(z) \otimes Z_{21}^{-1}(z)$$

$$O_1(\mu) = Z_{11} O_1^{(0)}, \quad Z_{11} = Z_m^{-1} = \frac{m_b(\mu)}{m_{b,0}}$$

$$H_2(z, \mu) = \int_0^1 dz' H_2^{(0)}(z') Z_{22}^{-1}(z', z)$$

$$O_2(z, \mu) = \int_0^1 dz' Z_{22}(z, z') O_2^{(0)}(z') + Z_{21}(z) O_1^{(0)}$$

$$\frac{\llbracket \bar{H}_2(z, \mu) \rrbracket}{z} = \int_0^{\boxed{\infty}} dz' \frac{\llbracket \bar{H}_2^{(0)}(z') \rrbracket}{z'} \llbracket Z_{22}^{-1}(z', z) \rrbracket$$

$$\llbracket O_2(z, \mu) \rrbracket = \int_0^{\boxed{\infty}} dz' \llbracket Z_{22}(z, z') \rrbracket \llbracket O_2^{(0)}(z') \rrbracket + \llbracket Z_{21}(z) \rrbracket O_1^{(0)}$$

$$H_3(\mu) = H_3^{(0)} Z_{33}^{-1}$$

$$J(\pm M_h \ell, \mu) = \int_0^{\boxed{\infty}} d\ell' Z_J(\pm M_h \ell, \pm M_h \ell') J^{(0)}(\pm M_h \ell')$$

$$S(w, \mu) = \int_0^{\boxed{\infty}} dw' Z_S(w, w'; \mu) S^{(0)}(w')$$

Bodwin, Ee, Lee, Wang, 2101.04872, 2107.07941


The red integral intervals are incompatible with subtraction/cutoff → **Mismatch**

$$Z_S(w, w') = \frac{w}{w'} Z_{33} \int_0^\infty \frac{dx}{x} Z_J^{-1}\left(\frac{M_h w'}{x \ell_+}, \frac{M_h w}{\ell_+}\right) Z_J^{-1}(-x M_h \ell_+, -M_h \ell_+),$$

determined by the
RG invariance of T3

Renormalization

Calculation of Z_{22} : UV divergences of matrix element of O_2



$$Z_{22}(z, z') = \left(1 + \frac{C_F \alpha_s}{4\pi\epsilon}\right) \delta(z - z')$$

Brodsky-Lepage Kernel


$$- \frac{C_F \alpha_s}{2\pi\epsilon} \frac{1}{z'(1-z')} \left[z(1-z') \frac{\theta(z' - z)}{z' - z} + z'(1-z) \frac{\theta(z - z')}{z - z'} \right]_+ + \mathcal{O}(\alpha_s^2)$$

$$\downarrow z, z' \rightarrow 0$$

$$[\![Z_{22}(z, z')]\!] = \left[1 - \frac{C_F \alpha_s}{4\pi\epsilon} (2 \ln z + 3)\right] \delta(z - z') - \frac{C_F \alpha_s}{2\pi\epsilon} z \left[\frac{\theta(z' - z)}{z'(z' - z)} + \frac{\theta(z - z')}{z(z - z')} \right]_+ + \mathcal{O}(\alpha_s^2)$$

$$Z_{33} [\![Z_{22}^{-1}(z, z')]\!] = M_h \frac{z}{z'} Z_J(z' M_h^2, z M_h^2) \quad \text{consistent with refactorization condition of H2}$$

Mismatch Terms in Renormalization



δT_2

$$- 2 \left(\int_0^1 dz \int_0^\infty dz' - \int_0^\infty dz \int_0^1 dz' \right) [\![H_2(z, \mu)]\!] [\![Z_{22}(z, z')]\!] [\langle O_2(z') \rangle]$$

$$+ (c \leftrightarrow \bar{c})$$

different integral intervals
but same integrands by
refactorization conditions

δT_3

$$H_3(\mu) \left[\int_0^{M_h} \frac{d\ell_-}{\ell_-} \int_0^{\sigma M_h} \frac{d\ell_+}{\ell_+} J(M_h \ell_-, \mu) J(-M_h \ell_+, \mu) S(\ell_+ \ell_-, \mu) \right.$$

$$\left. - \int_0^{M_h} \frac{d\ell_-}{\ell_-} \int_0^{\sigma M_h} \frac{d\ell_+}{\ell_+} J^{(0)}(M_h \ell_-) J^{(0)}(-M_h \ell_+) S^{(0)}(\ell_+ \ell_-) \right]$$

integrals over the entire planes
are scaleless

Sum of mismatch can be rearranged
to two integrals purely in hard region

Renormalized Matching Coefficients

Final renormalized H_1 :

$$\begin{aligned}
 H_1(\mu) &= \left(H_1^{(0)} + \Delta H_1^{(0)} - \delta H_1^{(0), \text{tot}} \right) Z_{11}^{-1} \\
 &+ 2 \lim_{\delta \rightarrow 0} \int_{\delta}^{1-\delta} dz \left[H_2^{(0)}(z) Z_{21}^{-1}(z) - \frac{\llbracket \bar{H}_2^{(0)}(z) \rrbracket}{z} \llbracket Z_{21}^{-1}(z) \rrbracket - \frac{\llbracket \bar{H}_2^{(0)}(\bar{z}) \rrbracket}{\bar{z}} \llbracket Z_{21}^{-1}(\bar{z}) \rrbracket \right] \\
 &\quad \text{Infinity-bin} \quad \text{Mismatch} \quad \text{Mixing from } O_2
 \end{aligned}$$

Final results at NLO:

$$H_1(\mu) = \frac{N_c \alpha_b}{\pi} \frac{y_b(\mu)}{\sqrt{2}} \left\{ -2 + \frac{C_F \alpha_s}{4\pi} \left[-\frac{\pi^2}{3} L_h^2 + (12 + 8\zeta_3) L_h - 36 - \frac{2\pi^2}{3} - \frac{11\pi^4}{45} \right] + \mathcal{O}(\alpha_s^2) \right\}$$

$$H_2(z, \mu) = \frac{y_b(\mu)}{\sqrt{2}} \frac{1}{z(1-z)} \left\{ 1 + \frac{C_F \alpha_s}{4\pi} \left[2L_h (\ln z + \ln(1-z)) + \ln^2 z + \ln^2(1-z) - 3 \right] + \mathcal{O}(\alpha_s^2) \right\},$$

$$\llbracket \bar{H}_2(z, \mu) \rrbracket = \frac{y_b(\mu)}{\sqrt{2}} \left[1 + \frac{C_F \alpha_s}{4\pi} (2L_h \ln z + \ln^2 z - 3) + \mathcal{O}(\alpha_s^2) \right],$$

$$H_3(\mu) = \frac{y_b(\mu)}{\sqrt{2}} \left[-1 + \frac{C_F \alpha_s}{4\pi} \left(L_h^2 + 2 - \frac{\pi^2}{6} \right) + \mathcal{O}(\alpha_s^2) \right]$$

$$L_h = \ln \frac{-M_h^2}{\mu^2}$$

Renormalized Matrix Elements

Final renormalized matrix elements at NLO:

$$\langle O_1(\mu) \rangle = m_b(\mu) g_\perp^{\mu\nu}$$

$$L_m = \ln \frac{m_b^2}{\mu^2} \quad \hat{w} = \frac{w}{m_b^2}$$

$$\begin{aligned} \langle O_2(z, \mu) \rangle &= \frac{N_c \alpha_b}{2\pi} m_b(\mu) g_\perp^{\mu\nu} \left\{ -L_m + \frac{C_F \alpha_s}{4\pi} \left[L_m^2 \left(\ln z + \ln(1-z) + 3 \right) \right. \right. \\ &\quad \left. \left. - L_m \left(\ln^2 z + \ln^2(1-z) - 4 \ln z \ln(1-z) + 11 - \frac{2\pi^2}{3} \right) + F(z) + F(1-z) \right] + \mathcal{O}(\alpha_s^2) \right\} \end{aligned}$$

$$\begin{aligned} [\![\langle O_2(z, \mu) \rangle]\!] &= \frac{N_c \alpha_b}{2\pi} m_b(\mu) g_\perp^{\mu\nu} \left\{ -L_m + \frac{C_F \alpha_s}{4\pi} \left[L_m^2 \left(\ln z + 3 \right) - L_m \left(\ln^2 z + 11 - \frac{2\pi^2}{3} \right) \right. \right. \\ &\quad \left. \left. + \frac{\ln^3 z}{6} - \frac{\ln z}{2} + 11 - \frac{\pi^2}{3} - 2\zeta_3 \right] + \mathcal{O}(\alpha_s^2) \right\} \end{aligned}$$

$$J(p^2, \mu) = 1 + \frac{C_F \alpha_s}{4\pi} \left[\ln^2 \left(\frac{-p^2 - i0}{\mu^2} \right) - 1 - \frac{\pi^2}{6} \right] + \mathcal{O}(\alpha_s^2)$$

$$S(w, \mu) = -\frac{N_c \alpha_b}{\pi} m_b(\mu) \left[S_a(w, \mu) \theta(w - m_b^2) + S_b(w, \mu) \theta(m_b^2 - w) \right]$$

$$S_a(w, \mu) = 1 + \frac{C_F \alpha_s}{4\pi} \left[-L_w^2 - 6L_w + 12 - \frac{\pi^2}{2} + 2 \text{Li}_2 \left(\frac{1}{\hat{w}} \right) - 4 \ln \left(1 - \frac{1}{\hat{w}} \right) \left(L_m + 1 + \ln \left(1 - \frac{1}{\hat{w}} \right) + \frac{3}{2} \ln \hat{w} \right) \right]$$

$$S_b(w, \mu) = \frac{C_F \alpha_s}{\pi} \ln(1 - \hat{w}) [L_m + \ln(1 - \hat{w})]$$

Renormalization-Group Equations

Hard functions

$$\frac{d}{d \ln \mu} H_1(\mu) = ?$$

$$\frac{d}{d \ln \mu} H_2(z, \mu) = \int_0^1 dz' H_2(z', \mu) \gamma_{22}(z', z)$$

$$\frac{d}{d \ln \mu} [\bar{H}_2(z, \mu)] = \int_0^\infty dz' [\bar{H}_2(z', \mu)] \frac{z}{z'} [\gamma_{22}(z', z)]$$

$$\frac{d}{d \ln \mu} H_3(\mu) = \gamma_{33} H_3(\mu)$$

Matrix elements

$$\frac{d}{d \ln \mu} \langle O_1(\mu) \rangle = -\gamma_{11} \langle O_1(\mu) \rangle$$

$$\begin{aligned} \frac{d}{d \ln \mu} \langle O_2(z, \mu) \rangle = & - \int_0^1 dz' \gamma_{22}(z, z') \langle O_2(z', \mu) \rangle \\ & - \gamma_{21}(z) \langle O_1(\mu) \rangle \end{aligned}$$

$$\begin{aligned} \frac{d}{d \ln \mu} [\langle O_2(z, \mu) \rangle] = & - \int_0^\infty dz' [\gamma_{22}(z, z')] [\langle O_2(z', \mu) \rangle] \\ & - [\gamma_{21}(z)] \langle O_1(\mu) \rangle \end{aligned}$$

$$\frac{d}{d \ln \mu} J(p^2, \mu) = - \int_0^\infty dx \gamma_J(p^2, xp^2) J(xp^2, \mu)$$

$$\frac{d}{d \ln \mu} S(w, \mu) = - \int_0^\infty dx \gamma_S(w, w/x) S(w/x, \mu)$$

- ★ RGE of H_1 is highly non-trivial because of the contributions from infinity-bin subtraction and mismatch in renormalization

Radiative Jet Function

Using the RG invariance of decay amplitude $B^- \rightarrow \gamma \ell^- \bar{\nu}$

$$\mathcal{M}(B^- \rightarrow \gamma \ell^- \bar{\nu}) \propto F_B(\mu) H(m_b, 2E_\gamma, \mu) \int_0^\infty \frac{d\omega}{\omega} J(-2E_\gamma \omega, \mu) \phi_+^B(\omega, \mu)$$

and the two-loop RGE of B meson LCDA, Braun, Ji, Manashov, '19

we can derive the two-loop anomalous dimension of jet function:

$$\begin{aligned} \gamma_J(p^2, xp^2; \mu) = & \left[\Gamma_{\text{cusp}}(\alpha_s) \ln \frac{-p^2}{\mu^2} - \gamma'(\alpha_s) \right] \delta(1-x) + \Gamma_{\text{cusp}}(\alpha_s) \Gamma(1, x) \\ & + C_F \left(\frac{\alpha_s}{2\pi} \right)^2 \frac{\theta(1-x)}{1-x} h(x) + \mathcal{O}(\alpha_s^3) \end{aligned}$$

Bell, Feldmann, Wang, Yip, '13

The solution of RGE of jet function at two loops is

jet function in dual space

$$\begin{aligned} J(p^2, \mu) = & \exp \left[-2S(\mu_j, \mu) - a_{\gamma'}(\mu_j, \mu) \right] \hat{\mathcal{J}}(\partial_\eta, \mu_j) \left(\frac{-p^2 e^{-2\gamma_E}}{\mu_j^2} \right)^{\eta + a_\Gamma(\mu_j, \mu)} \frac{\Gamma(1 - \eta - a_\Gamma(\mu_j, \mu))}{\Gamma(1 + \eta + a_\Gamma(\mu_j, \mu))} \\ & \times \exp \left[C_F \int_{\alpha_s(\mu_j)}^{\alpha_s(\mu)} \frac{d\alpha}{2\pi} \left[\mathcal{H}(\eta + a_\Gamma(\mu_j, \mu_\alpha)) + \mathcal{O}(\alpha) \right] \right] \Big|_{\eta=0} \end{aligned}$$

ZLL, Neubert, 2003.03393 (JHEP)

Soft-Quark Soft Function

RGE of soft-quark soft function: ZLL, Mecaj, Neubert, Wang, Fleming, 2005.03013 (JHEP)

$$\frac{d}{d \ln \mu} S(w, \mu) = - \int_0^\infty dw' \gamma_S(w, w'; \mu) S(w', \mu)$$

γ_S can be determined by RG invariance of T_3

$$\begin{aligned} \gamma_S(w, w'; \mu) = & - \left[\Gamma_{\text{cusp}}(\alpha_s) \ln \frac{w}{\mu^2} - \gamma_s(\alpha_s) \right] \delta(w - w') - 2\Gamma_{\text{cusp}}(\alpha_s) w \Gamma(w, w') \\ & - 2C_F \left(\frac{\alpha_s}{2\pi} \right)^2 \frac{w \theta(w' - w)}{w'(w' - w)} h\left(\frac{w}{w'}\right) + \mathcal{O}(\alpha_s^3) \end{aligned}$$

The solution of RGE gives:

$$S(w, \mu) = U_S(w; \mu, \mu_s) \left[\int_0^\infty \frac{dx}{x} S(w/x, \mu_s) G_{4,4}^{2,2} \left(\begin{matrix} -a, -a, 1-a, 1-a \\ 1, 1, 0, 0 \end{matrix} \middle| \frac{1}{x} \right) \right. \quad \text{Meijer G-Function} \\ \left. - m_b \frac{C_F \alpha_s(\mu_s)}{\pi} \int_0^1 \frac{dx}{1-x} \frac{h(x)}{\beta_0} \frac{r^{1+\frac{2C_F}{\beta_0} \ln x} - 1}{1 + \frac{2C_F}{\beta_0} \ln x} G_{4,4}^{2,2} \left(\begin{matrix} -a, -a, 1-a, 1-a \\ 0, 1, 0, 0 \end{matrix} \middle| \frac{xm_b^2}{w} \right) + \mathcal{O}(\alpha_s^2) \right]$$

Anomalous Dimensions

Anomalous dimensions:

$$\gamma_{ij} = \left(2\alpha_b \frac{\partial}{\partial \alpha_b} + 2\alpha_s \frac{\partial}{\partial \alpha_s} \right) Z_{ij}^{(1)}$$

$$\gamma_{11} = \frac{3C_F \alpha_s}{2\pi}$$

$$\gamma_{21}(z) = -\frac{N_c \alpha_b}{\pi} \left\{ 1 + \frac{C_F \alpha_s}{4\pi} \left[\ln^2 z + \ln^2(1-z) - 4 \ln z \ln(1-z) + 11 - \frac{2\pi^2}{3} \right] \right\}$$

$$[\![\gamma_{21}(z)]\!] = -\frac{N_c \alpha_b}{\pi} \left\{ 1 + \frac{C_F \alpha_s}{4\pi} \left(\ln^2 z + 11 - \frac{2\pi^2}{3} \right) \right\}$$

$$\gamma_{22}(z, z') = -\frac{C_F \alpha_s}{\pi} \left\{ \left[\ln z + \ln(1-z) + \frac{3}{2} \right] \delta(z - z') + z(1-z) \left[\frac{1}{z'(1-z)} \frac{\theta(z' - z)}{z' - z} + \frac{1}{z(1-z')} \frac{\theta(z - z')}{z - z'} \right]_+ \right\}$$

$$[\![\gamma_{22}(z, z')]\!] = -\frac{C_F \alpha_s}{\pi} \left\{ \left(\ln z + \frac{3}{2} \right) \delta(z - z') + z \left[\frac{\theta(z' - z)}{z'(z' - z)} + \frac{\theta(z - z')}{z(z - z')} \right]_+ \right\}$$

$$\gamma_{33} = \frac{C_F \alpha_s}{\pi} \left(L_h - \frac{3}{2} \right)$$

Determine RGE of H_1

RGE of H_1 has general formula:

$$\begin{aligned} \frac{d H_1(\mu)}{d \ln \mu} = & H_1(\mu) \gamma_{11}(\mu) + 2 \int_0^1 dz \left(H_2(z, \mu) \gamma_{21}(z, \mu) - \frac{\llbracket \bar{H}_2(z, \mu) \rrbracket}{z} \llbracket \gamma_{21}(z, \mu) \rrbracket - \frac{\llbracket \bar{H}_2(z, \mu) \rrbracket}{z} \llbracket \gamma_{21}(z, \mu) \rrbracket \right) \\ & + H_3(\mu) \gamma_{31}(\mu) \end{aligned}$$

RGE be determined by RG invariance:

$$\frac{d}{d \ln \mu} \mathcal{M}_b = 0 = \left[\left(\frac{d}{d \ln \mu} - \gamma_{11} \right) H_1(\mu) \right] \langle O_1(\mu) \rangle + \frac{d}{d \ln \mu} T_2(\mu) + \frac{d}{d \ln \mu} T_3(\mu)$$

The unknown quantity can be expressed as

non-zero due to cutoff

$$H_3(\mu) \gamma_{31}(\mu) \langle O_1(\mu) \rangle = -\frac{dT_3(\mu)}{d \ln \mu} + 4 \left(\int_0^1 dz \int_0^\infty dz' - \int_0^\infty dz \int_0^1 dz' \right) \frac{\llbracket \bar{H}_2(z', \mu) \rrbracket}{z'} \llbracket \gamma_{22}(z', z, \mu) \rrbracket \llbracket \langle O_2(z, \mu) \rangle \rrbracket$$

mismatch in T_2

each of them depends on hard and soft scales,
but their combination is purely hard

Determine RGE of H_1

With the two refactorization conditions:

$$\begin{aligned}
 & 2 \left(\int_0^1 dz \int_0^\infty dz' - \int_0^\infty dz \int_0^1 dz' \right) \frac{[\bar{H}_2(z', \mu)]}{z'} [\gamma_{22}(z', z, \mu)] [\langle O_2(z, \mu) \rangle] \\
 & = H_3(\mu) \int_0^\infty dx K(x) \left\{ \int_{M_h}^{M_h/x} \frac{d\ell_-}{\ell_-} \int_0^\infty \frac{d\ell_+}{\ell_+} J(x M_h \ell_-, \mu; \epsilon) J(-M_h \ell_+, \mu; \epsilon) S(\ell_+ \ell_-, \mu; \epsilon) \right. \\
 & \quad \left. - 2 \int_{M_h}^{M_h/x} \frac{d\ell_-}{\ell_-} J(x M_h \ell_-, \mu; \epsilon) [Z_{21}(\ell_- / M_h, \mu)] Z_{11}^{-1}(\mu) \langle O_1(\mu) \rangle \right\}
 \end{aligned}$$

To regulate the endpoint divergences, one need to keep the full dependence on ϵ in renormalized jet and soft functions before performing convolution

On the other hand:

$$\frac{dT_3(\mu)}{d \ln \mu} = H_3(\mu) \int_0^\infty dx K(x, \mu) \int_{M_h}^{M_h/x} \frac{d\ell_-}{\ell_-} \int_0^{\sigma M_h} \frac{d\ell_+}{\ell_+} J(x M_h \ell_-, \mu) J(-M_h \ell_+, \mu) S(\ell_+ \ell_-, \mu) + (\ell_+ \leftrightarrow \ell_-)$$

with $K(x, \mu) = \Gamma_{\text{cusp}}(\alpha_s) \Gamma(1, x) + C_F \left(\frac{\alpha_s}{2\pi} \right)^2 \frac{\theta(1-x)}{1-x} h(x) + \mathcal{O}(\alpha_s^3)$

Combination of the two mismatch terms gives an integral in purely hard region

RG Evolution


Last piece of anomalous dimensions:

$$\gamma_{31}(\mu) = \frac{N_c \alpha_b}{\pi} \sum_{n=0}^{\infty} \gamma_{31}^{(n)} \left(\frac{\alpha_s}{4\pi} \right)^{n+1}$$

$$\gamma_{31}^{(0)} = 16\zeta_3 C_F,$$

$$\begin{aligned} \gamma_{31}^{(1)} = & C_F^2 \left[-32\zeta_3 L_h^2 + \left(\frac{8\pi^4}{15} - 48\zeta_3 \right) L_h \right. \\ & + \frac{8\pi^2\zeta_3}{3} - 32\zeta_5 + \frac{4\pi^4}{5} + 168\zeta_3 \left. \right] \\ & + C_F C_A \left(-\frac{176\zeta_3}{3} L_h - \frac{16\pi^2\zeta_3}{3} - \frac{44\pi^4}{45} + \frac{1072\zeta_3}{9} \right) \\ & + C_F n_f T_F \left(\frac{64\zeta_3}{3} L_h + \frac{16\pi^4}{45} - \frac{320\zeta_3}{9} \right) \end{aligned}$$

Dynamical scale choice in T_3



- Double log term in anomalous dimension complicates the scale evolution of H_1
- Evolving the matrix elements up to hard scale is accessible
- Dynamical scale setting is necessary to adapt to the convolution

Resummation at NLL

Evolved quantities in T3 with dynamical scale setting

$$H_{3,\text{LO}}(\mu) = -\frac{y_b(M_h)}{\sqrt{2}} e^{2S_\Gamma(-M_h^2, \mu^2) - 2a_{\gamma_q}(-M_h^2, \mu^2)},$$

$$J_{\text{LO}}(M_h \ell_-, \mu) = e^{-2S_\Gamma(-M_h \ell_-, \mu^2) - a_{\gamma'}(-M_h \ell_-, \mu^2) - 2\gamma_E a_-} \frac{\Gamma(1 - a_-)}{\Gamma(1 + a_-)},$$

$$J_{\text{LO}}(-M_h \ell_+, \mu) = e^{-2S_\Gamma(M_h \ell_+, \mu^2) - a_{\gamma'}(M_h \ell_+, \mu^2) - 2\gamma_E a_+} \frac{\Gamma(1 - a_+)}{\Gamma(1 + a_+)},$$

$$S_{\text{LO}}(\ell_+ \ell_-, \mu) = -\frac{N_c \alpha_b}{\pi} m_b e^{2S_\Gamma(\ell_+ \ell_-, \mu^2) + a_{\gamma_s}(\ell_+ \ell_-, \mu^2) + 4\gamma_E a_s}$$

$$\times \left(1 - \frac{\alpha_s(\ell_+ \ell_-)}{8\pi} \gamma_{m,0} \ln \frac{m_b^2}{\ell_+ \ell_-}\right) G_{4,4}^{2,2} \left(\begin{matrix} -a_s, -a_s, 1 - a_s, 1 - a_s \\ 0, 1, 0, 0 \end{matrix} \middle| \frac{m_b^2}{w} \right)$$

$$\begin{aligned} S_\Gamma(\nu, \mu) &= \frac{\alpha_s(\nu)}{4\pi} \left(-\frac{\Gamma_0}{8} \ln^2 \frac{\mu^2}{\nu^2} \right) \\ &\quad + \left(\frac{\alpha_s(\nu)}{4\pi} \right)^2 \left(\frac{\beta_0 \Gamma_0}{12} \ln^3 \frac{\mu^2}{\nu^2} + \dots \right) + \dots \\ a_\Gamma(\nu, \mu) &= \frac{\alpha_s(\nu)}{4\pi} \left(-\frac{\Gamma_0}{2} \ln \frac{\mu^2}{\nu^2} \right) + \dots \\ \alpha_s(\mu) &= \alpha_s(\nu) \left(1 - \frac{\alpha_s(\nu)}{4\pi} \beta_0 \ln \frac{\mu^2}{\nu^2} + \dots \right) \end{aligned}$$

$$a_- = a_\Gamma(-M_h \ell_-, \mu^2)$$

$$a_+ = a_\Gamma(M_h \ell_+, \mu^2)$$

$$a_s = a_\Gamma(\ell_+ \ell_-, \mu^2)$$

NLL Resummation for decay amplitude

$$\theta(\ell_+ \ell_- - m_b^2) \frac{\Gamma^2(1 + a_s)}{\Gamma^2(1 - a_s)} + \mathcal{O}\left(\frac{m_b^2}{\ell_+ \ell_-}\right)$$

$$\begin{aligned} \mathcal{M}_b^{\text{NLL}} &= \frac{N_c \alpha_b}{\pi} \frac{y_b(M_h)}{\sqrt{2}} m_b \varepsilon_\perp^*(k_1) \cdot \varepsilon_\perp^*(k_2) \frac{L^2}{2} \left\{ \sum_{n=0}^{\infty} \frac{2\Gamma(n+1)}{\Gamma(2n+3)} \left(-\frac{C_F \alpha_s(M_h)}{2\pi} L^2 \right)^n \right. \\ &\quad \left. - \frac{1}{L} \sum_{n=1}^{\infty} \frac{\Gamma(n+1)}{\Gamma(2n+2)} \left(-\frac{C_F \alpha_s(M_h)}{2\pi} L^2 \right)^n \left[3 - \beta_0 \frac{\alpha_s(M_h)}{2\pi} L^2 \frac{n(n+1)}{(2n+2)(2n+3)} \right] \right\} \end{aligned}$$

Logarithms at Three Loops

Solving the RGEs iteratively in perturbation theory, one can predict the large logarithms in three-loop decay amplitude down to $\alpha\alpha_s^2 L^3$

$$\begin{aligned} \mathcal{M}_b = & \frac{N_c \alpha_b}{\pi} \frac{m_b^2}{v} \varepsilon_{\perp}^*(k_1) \cdot \varepsilon_{\perp}^*(k_2) \\ & \times \left\{ \frac{L^2}{2} - 2 + \frac{C_F \alpha_s(\hat{\mu}_h)}{4\pi} \left[-\frac{L^4}{12} - L^3 - \frac{2\pi^2}{3} L^2 + \left(12 + \frac{2\pi^2}{3} + 16\zeta_3 \right) L - 20 + 4\zeta_3 - \frac{\pi^4}{5} \right] \right. \\ & \left. + C_F \left(\frac{\alpha_s(\hat{\mu}_h)}{4\pi} \right)^2 \left[\frac{C_F}{90} L^6 + \left(\frac{C_F}{10} - \frac{\beta_0}{30} \right) L^5 + d_4^{\text{OS}} L^4 + d_3^{\text{OS}} L^3 + \dots \right] \right\} \end{aligned}$$

agree with numerical result of Niggetto, 2020

Strong evidence for the correctness of our factorization !!!


with

$$d_4^{\text{OS}} = \left(\frac{3}{2} + \frac{\pi^2}{18} \right) C_F + \left(-\frac{91}{27} + \frac{\pi^2}{36} \right) C_A + \frac{32}{27} T_F n_f ,$$

$$d_3^{\text{OS}} = \left(-\frac{1}{2} + \frac{7\pi^2}{9} + \frac{20}{3} \zeta_3 \right) C_F + \left(-\frac{199}{18} - \frac{22\pi^2}{27} - 4\zeta_3 \right) C_A + \left(\frac{22}{9} + \frac{8\pi^2}{27} \right) T_F n_f$$

Is conventional QCD approach valid beyond leading logarithmic accuracy?

Resummation beyond NLL



Resummation of T_2

Employ Gegenbauer Polynomial: $O_2(z, \mu) = z(1-z) \sum_{n=0}^{\infty} a_n(\mu) C_n^{(3/2)}(2z-1)$

The convolution can be written as: $\int_0^1 dz H_2(z, \mu) O_2(z, \mu) = \sum_{n=0}^{\infty} a_{2n}(\mu) h_{2n}(\mu)$

Leads to local RGE: $\frac{d}{d \ln \mu} a_{2n}(\mu) = -\gamma_{21,2n}(\mu) O_1(\mu) - \gamma_{22,2n}(\mu) a_{2n}(\mu)$

$$\gamma_{21,2n}(\mu) = \left(-\frac{N_c \alpha_b}{\pi}\right) \frac{2(4n+3)}{(2n+1)(n+1)} \left\{ 1 + \frac{C_F \alpha_s}{4\pi} \left[4H_{2n+1}^2 - 2\psi^{(1)}(n+1) + 2\psi^{(1)}\left(n+\frac{3}{2}\right) + 11 - \frac{2\pi^2}{3} \right] \right\}$$

$$\gamma_{22,2n}(\mu) = \frac{C_F \alpha_s}{4\pi} (8H_{2n+1} - 6)$$

The solution of RGE:

$$\mu = M_h \quad \mu_0 = m_b$$

$$a_{2n}(\mu) = -\frac{N_c \alpha_b}{2\pi} m_b(\mu_0) \frac{2(4n+3)}{(2n+1)(n+1)} \times \left\{ \left(\frac{\alpha_s}{4\pi}\right)^{-1} [C_F(6 - 4H_{2n+1}) - \beta_0]^{-1} \left[\left(\frac{\alpha_s(\mu)}{\alpha_s(\mu_0)}\right)^{\frac{\gamma_{11}^0}{2\beta_0}} - \left(\frac{\alpha_s(\mu)}{\alpha_s(\mu_0)}\right)^{\frac{\gamma_{22,2n}^0}{2\beta_0} + 1} \right] \right\}$$

Divergent
Convergent

Resummation of T_2

Transform to the dual space:

$$[\tilde{O}_2(z, \mu)] = \int_0^\infty \frac{dx}{\sqrt{x}} \mathbf{J}_1(2\sqrt{x}) [O_2(xz, \mu)]$$

Leading to local RGE:

$$\frac{d}{d \ln \mu} [\tilde{O}_2(z, \mu)] = -[\tilde{\gamma}_{22}(z, \mu)] [\tilde{O}_2(z, \mu)] - [\tilde{\gamma}_{21}(z, \mu)] O_1(\mu)$$

$$[\tilde{\gamma}_{21}(z, \mu)] = -\frac{N_c \alpha_b}{\pi} \left[1 + \frac{C_F \alpha_s}{4\pi} \left(\ln^2(z e^{-2\gamma_E}) + 11 - \frac{2\pi^2}{3} \right) \right]$$

$$[\tilde{\gamma}_{22}(z, \mu)] = \frac{C_F \alpha_s}{4\pi} (-4 \ln z + 8\gamma_E - 6)$$

The solution of RGE:

$$[O_2(z, \mu)] = \left(-\frac{N_c \alpha_b}{2\pi} \right) m_b(\mu_0) \left(\frac{\alpha_s}{4\pi} \right)^{-1} [C_F(2\partial_\eta + 6) - \beta_0]^{-1}$$

$$\times \left[\left(\frac{\alpha_s(\mu)}{\alpha_s(\mu_0)} \right)^{\frac{\gamma_{11}^0}{2\beta_0}} (ze^{-2\gamma_E})^\eta \frac{\Gamma(1-\eta)}{\Gamma(1+\eta)} - \left(\frac{\alpha_s(\mu)}{\alpha_s(\mu_0)} \right)^{1-\frac{3C_F}{\beta_0}} (ze^{-2\gamma_E})^{-a+\eta} \frac{\Gamma(1-\eta+a)}{\Gamma(1+\eta-a)} \right] \Big|_{\eta=0}$$

Divergent
Convergent

$$\frac{1}{\partial_\eta - x} = (1 + x\partial_\eta^{-1} + x^2\partial_\eta^{-2} + x^3\partial_\eta^{-3} + \dots) \partial_\eta^{-1}$$

Conclusions

- First renormalized factorization theorem at subleading power (with endpoint divergences) is established in SCET
- Two refactorization conditions are proved at operator level
- Nature cutoffs arise from subtractions and rearrangement of endpoint singularities
- Contributions from mismatch between renormalization and convolution with cutoffs are proved to be purely hard
- Operators mix under renormalization

Thanks for your attention!