

CLASSICAL AND QUANTUM ASPECTS OF THE STRING DOUBLE SIGMA MODEL

Franco Pezzella

INFN - Naples Division

STRINGY GEOMETRY - Mainz - September, 15th 2015

BASED ON:

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- F. P., [arXiv:1503.01709](#).
- L. De Angelis, G. Gionti, R. Marotta and F. P., [JHEP 1404 \(2014\) 171](#), [arXiv:1312.7367](#).
- F. Rennecke, [JHEP 1410 \(2014\) 69](#) .
- M. Duff, [Nucl. Phys. B335 \(1990\) 610](#).
- A. A. Tseytlin, [Phys.Lett. B242 \(1990\) 163-174](#) ; [Nucl.Phys. B350 \(1991\) 395-440](#).
- R. Floreanini and R. Jackiw, [Phys. Rev. Lett. 59 \(1987\) 1873](#).
- C. Hull, [JHEP 0510 \(2005\) 065](#), [hep-th/0406102](#) .

PLAN OF THE TALK

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

1 INTRODUCTION AND MOTIVATION

PLAN OF THE TALK

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

1 INTRODUCTION AND MOTIVATION

2 HODGE-DUAL SYMMETRIC FREE SCALAR FIELDS IN 2D

PLAN OF THE TALK

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- 1 INTRODUCTION AND MOTIVATION
- 2 HODGE-DUAL SYMMETRIC FREE SCALAR FIELDS IN 2D
- 3 DOUBLE SIGMA MODEL (CLOSED STRINGS)

PLAN OF THE TALK

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- 1 INTRODUCTION AND MOTIVATION
- 2 HODGE-DUAL SYMMETRIC FREE SCALAR FIELDS IN 2D
- 3 DOUBLE SIGMA MODEL (CLOSED STRINGS)
- 4 DUALITY SYMMETRIC FREE CLOSED STRINGS

PLAN OF THE TALK

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- 1 INTRODUCTION AND MOTIVATION
- 2 HODGE-DUAL SYMMETRIC FREE SCALAR FIELDS IN 2D
- 3 DOUBLE SIGMA MODEL (CLOSED STRINGS)
- 4 DUALITY SYMMETRIC FREE CLOSED STRINGS
- 5 QUANTIZATION OF THE DOUBLE STRING MODEL

PLAN OF THE TALK

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- 1 INTRODUCTION AND MOTIVATION
- 2 HODGE-DUAL SYMMETRIC FREE SCALAR FIELDS IN 2D
- 3 DOUBLE SIGMA MODEL (CLOSED STRINGS)
- 4 DUALITY SYMMETRIC FREE CLOSED STRINGS
- 5 QUANTIZATION OF THE DOUBLE STRING MODEL
- 6 CONCLUSION AND PERSPECTIVES

REMINDING T-DUALITY

- **T-duality** is an old subject in string theory. It implies that in many cases two different geometries for the extra-dimensions are physically equivalent.

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

REMINDING T-DUALITY

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- **T-duality** is an old subject in string theory. It implies that in many cases two different geometries for the extra-dimensions are physically equivalent.
- T-duality is a discrete symmetry. It implies that string physics at a very small scale cannot be distinguished from the one at a large scale. It is also a clear indication that **ordinary geometric concepts can break down in string theory at the string scale.**

REMINING T-DUALITY

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- **T-duality** is an old subject in string theory. It implies that in many cases two different geometries for the extra-dimensions are physically equivalent.
- T-duality is a discrete symmetry. It implies that string physics at a very small scale cannot be distinguished from the one at a large scale. It is also a clear indication that **ordinary geometric concepts can break down in string theory at the string scale.**
- In the simplest case of **circular compactifications**, T-duality is encoded, for bosonic closed strings, in the simultaneous transformations $R \leftrightarrow \alpha'/R$ and $p_a \leftrightarrow w^a/\alpha'$ under which $X^a = X_L^a + X_R^a \leftrightarrow \tilde{X}_a \equiv X_L^a - X_R^a$, with w^a playing the role of momentum mode for \tilde{X}_a . These transformations leave the **mass spectrum** invariant.

REMINING T-DUALITY

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- **T-duality** is an old subject in string theory. It implies that in many cases two different geometries for the extra-dimensions are physically equivalent.
- T-duality is a discrete symmetry. It implies that string physics at a very small scale cannot be distinguished from the one at a large scale. It is also a clear indication that **ordinary geometric concepts can break down in string theory at the string scale.**
- In the simplest case of **circular compactifications**, T-duality is encoded, for bosonic closed strings, in the simultaneous transformations $R \leftrightarrow \alpha'/R$ and $p_a \leftrightarrow w^a/\alpha'$ under which $X^a = X_L^a + X_R^a \leftrightarrow \tilde{X}_a \equiv X_L^a - X_R^a$, with w^a playing the role of momentum mode for \tilde{X}_a . These transformations leave the **mass spectrum** invariant.
- In **toroidal compactifications** (with constant backgrounds $G_{\mu\nu}$ and $B_{\mu\nu}$) T-duality is described by $O(D, D; \mathbb{Z})$ transformations.

O(D,D) DUALITY IN STRING THEORY

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- Already on the classical level the indefinite orthogonal group $O(D, D; \mathbb{R})$ appears naturally in the Hamiltonian description of the usual bosonic string model.
- With $*$ the Hodge operator with respect to $h = \text{diag}(-1, 1)$, the action is:

$$S[X; G, B] = \frac{T}{2} \int [G_{ab}(X) dX^a \wedge *dX^b + B_{ab}(X) dX^a \wedge dX^b]$$

O(D,D) DUALITY IN STRING THEORY

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- Already on the classical level the indefinite orthogonal group $O(D, D; \mathbb{R})$ appears naturally in the Hamiltonian description of the usual bosonic string model.
- With $*$ the Hodge operator with respect to $h = \text{diag}(-1, 1)$, the action is:

$$S[X; G, B] = \frac{T}{2} \int [G_{ab}(X) dX^a \wedge *dX^b + B_{ab}(X) dX^a \wedge dX^b]$$

- Varying S with respect to X^a yields the equation of motion:

$$d * dX^a + \Gamma^a_{bc} dX^b \wedge *dX^c = \frac{1}{2} G^{am} H_{mbc} dX^b \wedge dX^c \quad (1)$$

with $H = dB$ and $\Gamma^a_{bc} = \frac{1}{2} G^{am} (\partial_b G_{mc} + \partial_c G_{mb} - \partial_m G_{bc})$ the coefficients of the Levi Civita connection.

O(D,D) DUALITY IN STRING THEORY

- The dynamics of the theory is determined by the equations of motion for the coordinates X^a accompanied with the constraints (in the conformal gauge):

$$G_{ab}(\dot{X}^a \dot{X}^b + X'^a X'^b) = 0 \quad G_{ab} \dot{X}^a X'^b = 0. \quad (2)$$

These derive from the vanishing of the energy-momentum tensor $T_{\alpha\beta} = 0$, i.e. from the equation of motion for a general world-sheet metric h .

O(D,D) DUALITY IN STRING THEORY

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The dynamics of the theory is determined by the equations of motion for the coordinates X^a accompanied with the constraints (in the conformal gauge):

$$G_{ab}(\dot{X}^a \dot{X}^b + X'^a X'^b) = 0 \quad G_{ab} \dot{X}^a X'^b = 0. \quad (2)$$

These derive from the vanishing of the energy-momentum tensor $T_{\alpha\beta} = 0$, i.e. from the equation of motion for a general world-sheet metric h .

- The Hamiltonian density can be determined from the Lagrangian density by performing a Legendre transformation with respect to the **canonical momentum**

$$P_a = \frac{\partial L}{\partial \dot{X}^a} = \frac{1}{2\pi\alpha'} \left(-G_{ab} \dot{X}^b + B_{ab} X'^b \right) \text{ and } \dot{X}^a.$$

O(D,D) DUALITY IN STRING THEORY

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The dynamics of the theory is determined by the equations of motion for the coordinates X^a accompanied with the constraints (in the conformal gauge):

$$G_{ab}(\dot{X}^a \dot{X}^b + X'^a X'^b) = 0 \quad G_{ab} \dot{X}^a X'^b = 0. \quad (2)$$

These derive from the vanishing of the energy-momentum tensor $T_{\alpha\beta} = 0$, i.e. from the equation of motion for a general world-sheet metric h .

- The Hamiltonian density can be determined from the Lagrangian density by performing a Legendre transformation with respect to the **canonical momentum**

$$P_a = \frac{\partial L}{\partial \dot{X}^a} = \frac{1}{2\pi\alpha'} \left(-G_{ab} \dot{X}^b + B_{ab} X'^b \right) \text{ and } \dot{X}^a.$$

- By virtue of the constraint $G_{ab} \dot{X}^a X'^b = 0$, the Hamiltonian density can also result from a Legendre transformation with respect to the **canonical winding**

$$W_a = \frac{\partial L}{\partial X'^a} = \frac{1}{2\pi\alpha'} \left(G_{ab} X'^b - B_{ab} \dot{X}^b \right) \text{ and } X'^a.$$

O(D,D) INVARIANCE OF THE STRING HAMILTONIAN DENSITY

- The Hamiltonian density can be written equivalently as:

$$\begin{aligned} H &= -\frac{1}{4\pi\alpha'} \left(\begin{array}{c} \partial_\sigma X \\ 2\pi\alpha' P \end{array} \right)^t \mathcal{M}(G, B) \left(\begin{array}{c} \partial_\sigma X \\ 2\pi\alpha' P \end{array} \right) \\ &= -\frac{1}{4\pi\alpha'} \left(\begin{array}{c} \partial_\tau X \\ -2\pi\alpha' W \end{array} \right)^t \mathcal{M}(G, B) \left(\begin{array}{c} \partial_\tau X \\ 2\pi\alpha' W \end{array} \right) \end{aligned}$$

where the *generalised metric* is introduced:

$$\mathcal{M}(G, B) = \begin{pmatrix} G - BG^{-1}B & BG^{-1} \\ -G^{-1}B & -G^{-1} \end{pmatrix} \quad (3)$$

- Defining the generalised vectors $A_P(X)$ and $A_W(X)$ in $TM \oplus T^*M$ with components

$$A_P(X) = \left(\begin{array}{c} \partial_\sigma X \\ 2\pi\alpha' P \end{array} \right) \quad A_W(X) = \left(\begin{array}{c} \partial_\tau X \\ 2\pi\alpha' W \end{array} \right)$$

one can see that the Hamiltonian density is proportional to the squared length of A_P and A_W as measured by the generalised metric \mathcal{M} .

CONSTRAINTS AND GENERALISED VECTORS

- In terms of the generalised vector A_P the constraints, i.e. the components of the energy-momentum tensor can be rewritten as:

$$A_P^t \mathcal{M} A_P = 0 \quad A_P^t \Omega A_P = 0. \quad (4)$$

The first constraint sets the Hamiltonian density to zero, hence the second constraint completely determines the dynamics and it is rewritten in terms of the matrix $\Omega = \begin{pmatrix} 0 & \mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix}$, i.e. the invariant metric of the group $O(D, D)$ defined by the $D \times D$ matrices T satisfying the condition $T^t \Omega T = \Omega$. In particular the generalised metric is an element of $O(D, D)$.

- All the admissible generalised vectors satisfying the constraints are related by $O(D, D)$ transformations, which implies a simultaneous inverse transformation of the generalised metric.

This, in turn, leaves the Hamiltonian density and the energy-momentum tensor invariant.

$O(D, D; R)$ IN THE PRESENCE OF CONSTANT BACKGROUNDS

CLASSICAL AND QUANTUM ASPECTS OF THE STRING DOUBLE SIGMA MODEL

FRANCO PEZZELLA

INTRODUCTION AND MOTIVATION

HODGE-DUAL SYMMETRIC FREE SCALAR FIELDS IN 2D

DOUBLE SIGMA MODEL (CLOSED STRINGS)

DUALITY SYMMETRIC FREE CLOSED STRINGS

QUANTIZATION OF THE DOUBLE STRING MODEL

CONCLUSION AND PERSPECTIVES

- In the presence of constant backgrounds (G, B) , the equations of motion for the string coordinates are a set of conservation laws on the world-sheet:

$$\partial_\alpha J_a^\alpha = 0 \rightarrow J_a^\alpha = G_{ab} \partial_\alpha X^b + \epsilon^{\alpha\beta} B_{ab} \partial_\beta X^b \quad (5)$$

- Locally, one can express such currents as:

$$G_{ab} \partial_\alpha X^b + \epsilon^{\alpha\beta} B_{ab} \equiv -\epsilon^{\alpha\beta} \partial_\beta \tilde{X}_a \rightarrow \text{dual coordinates}$$

in terms of which the action S can be rewritten as:

$$S[X; G, B] = \frac{T}{2} \int \left[\tilde{G}_{ab} d\tilde{X}^a \wedge *d\tilde{X}^b + \tilde{B}_{ab}(X) d\tilde{X}^a \wedge d\tilde{X}^b \right]$$

with $\tilde{G} = G - BG^{-1}B$ and $\tilde{B} = -\tilde{G}B^{-1}G$.

- The equations of motion for the coordinates $\chi = (X, \tilde{X})$ can be combined into a single equation $O(D, D)$ -invariant:

$$M \partial_\alpha \chi = \Omega \epsilon_{\alpha\beta} \partial^\beta \chi \quad (6)$$

- For $B = 0$, the equations of motion become the duality conditions: $\partial_\alpha X^a = -\epsilon_{\alpha\beta} \partial^\beta \tilde{X}^a$.

$O(D, D; \mathbb{R}) \rightarrow O(D, D; \mathbb{Z})$

- If the closed string coordinates are defined on a compact target manifold, the dual coordinates will satisfy the same periodicity conditions and then T-duality maps two theories of the same type into one another \rightarrow **exact symmetry**.
- For closed strings, toroidal compactification means:

$$X^a(\sigma, \tau) \equiv X^a(\sigma + \pi, \tau) + 2\pi L^a \quad L^a = \sum_{i=1}^d w_i R_i e_i^a \quad (7)$$

with w_i being the winding numbers and e_i^a vector basis on T^d .

- In the compact space $O(D, D; \mathbb{R}) \rightarrow O(D, D; \mathbb{Z})$. The latter becomes the T-duality group of the toroidal compactification. For closed strings on compactified dimensions, this group becomes a *symmetry* not only of the mass spectrum and the vacuum partition function but also of the scattering amplitudes.

T-DUAL INVARIANT BOSONIC STRING FORMULATION

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The presence of the $O(D, D)$ symmetry suggests to extend the standard formulation of String Theory, based on the Polyakov action, by introducing this symmetry at the level of the world-sheet sigma-model. It would be interesting, therefore, looking for a **manifestly $O(D, D)$ -dual invariant formulation of the string theory**.
- The introduction of *both* the coordinates X^a *and* the dual ones \tilde{X}_a is required. Such formulation is based on a **doubling** of the string coordinates in the target space.

DOUBLING COORDINATES: MOTIVATION

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The main goal of this new action would be **to explore more closely aspects of stringy geometry** and, in particular, of string gravity. In fact, if interested in writing down the complete effective field theory of such generalised sigma-model, one should consider, correspondingly to the introduction of X^a and \tilde{X}_a , a dependence of the fields associated with string states on such coordinates. In this way, **double string effective field theory** becomes a **double field theory**.

DOUBLING COORDINATES: MOTIVATION

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The main goal of this new action would be **to explore more closely aspects of stringy geometry** and, in particular, of string gravity. In fact, if interested in writing down the complete effective field theory of such generalised sigma-model, one should consider, correspondingly to the introduction of X^a and \tilde{X}_a , a dependence of the fields associated with string states on such coordinates. In this way, **double string effective field theory** becomes a **double field theory**.
- What the well-known **effective gravitational action of a closed string**

$$S = \int dX \sqrt{G} e^{-2\phi} \left[R + 4(\partial\phi)^2 - \frac{1}{12} H_{\mu\nu\rho} H^{\mu\nu\rho} \right]$$

becomes when G , B and ϕ are dependent on X^a and \tilde{X}_a ? Which symmetries and what properties would it have? This could shed light on aspects of string gravity unexplored thus far.

DOUBLING COORDINATES: MOTIVATION

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The main goal of this new action would be **to explore more closely aspects of stringy geometry** and, in particular, of string gravity. In fact, if interested in writing down the complete effective field theory of such generalised sigma-model, one should consider, correspondingly to the introduction of X^a and \tilde{X}_a , a dependence of the fields associated with string states on such coordinates. In this way, **double string effective field theory** becomes a **double field theory**.

- What the well-known **effective gravitational action of a closed string**

$$S = \int dX \sqrt{G} e^{-2\phi} \left[R + 4(\partial\phi)^2 - \frac{1}{12} H_{\mu\nu\rho} H^{\mu\nu\rho} \right]$$

becomes when G , B and ϕ are dependent on X^a and \tilde{X}_a ? Which symmetries and what properties would it have? This could shed light on aspects of string gravity unexplored thus far.

- How the string theory would look like when the T-duality is manifested in the sigma-model Lagrangian density?

HODGE-DUALITY SYMMETRY FOR 2D SCALAR FIELDS

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The usual Lagrangian of a 2D scalar field ϕ

$$\mathcal{L} = -\frac{1}{2}\partial_\alpha\phi\partial^\alpha\phi = \frac{1}{2}\eta^{\alpha\beta}\partial_\alpha\phi\partial_\beta\phi = \frac{1}{2}\dot{\phi}^2 - \frac{1}{2}\phi'^2$$

can be rewritten in a manifestly invariant form under $\phi \leftrightarrow \tilde{\phi}$, its Hodge dual defined by $\partial_\alpha\tilde{\phi} = -\epsilon_{\alpha\beta}\partial^\beta\phi$ ($\epsilon^{01} = 1$).

HODGE-DUALITY SYMMETRY FOR 2D SCALAR FIELDS

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The usual Lagrangian of a 2D scalar field ϕ

$$\mathcal{L} = -\frac{1}{2}\partial_\alpha\phi\partial^\alpha\phi = \frac{1}{2}\eta^{\alpha\beta}\partial_\alpha\phi\partial_\beta\phi = \frac{1}{2}\dot{\phi}^2 - \frac{1}{2}\phi'^2$$

can be rewritten in a manifestly invariant form under $\phi \leftrightarrow \tilde{\phi}$, its Hodge dual defined by $\partial_\alpha\tilde{\phi} = -\epsilon_{\alpha\beta}\partial^\beta\phi$ ($\epsilon^{01} = 1$).

- Two steps are necessary.

HODGE-DUALITY SYMMETRY FOR 2D SCALAR FIELDS

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The usual Lagrangian of a 2D scalar field ϕ

$$\mathcal{L} = -\frac{1}{2}\partial_\alpha\phi\partial^\alpha\phi = \frac{1}{2}\eta^{\alpha\beta}\partial_\alpha\phi\partial_\beta\phi = \frac{1}{2}\dot{\phi}^2 - \frac{1}{2}\phi'^2$$

can be rewritten in a manifestly invariant form under $\phi \leftrightarrow \tilde{\phi}$, its Hodge dual defined by $\partial_\alpha\tilde{\phi} = -\epsilon_{\alpha\beta}\partial^\beta\phi$ ($\epsilon^{01} = 1$).

- Two steps are necessary.
- The first consists in rewriting \mathcal{L} in a first order form, after introducing an auxiliary field p whose equation of motion reproduces $p = \dot{\phi}$.

HODGE-DUALITY SYMMETRY FOR 2D SCALAR FIELDS

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The usual Lagrangian of a 2D scalar field ϕ

$$\mathcal{L} = -\frac{1}{2}\partial_\alpha\phi\partial^\alpha\phi = \frac{1}{2}\eta^{\alpha\beta}\partial_\alpha\phi\partial_\beta\phi = \frac{1}{2}\dot{\phi}^2 - \frac{1}{2}\phi'^2$$

can be rewritten in a manifestly invariant form under $\phi \leftrightarrow \tilde{\phi}$, its Hodge dual defined by $\partial_\alpha\tilde{\phi} = -\epsilon_{\alpha\beta}\partial^\beta\phi$ ($\epsilon^{01} = 1$).

- Two steps are necessary.
- The first consists in rewriting \mathcal{L} in a first order form, after introducing an auxiliary field p whose equation of motion reproduces $p = \dot{\phi}$.
- The second consists in trading p for the new field $\tilde{\phi}$ defined through $p \equiv \tilde{\phi}'$. It is easy to see that this procedure leads to the following symmetric Lagrangian:

$$\mathcal{L}_{sym} = \frac{1}{2} \left[\dot{\phi}\tilde{\phi}' + \phi'\tilde{\phi} - \phi'^2 - \tilde{\phi}'^2 \right]$$

- The manifest Lorentz invariance has disappeared, but it holds on-shell.

FREE SCALAR FIELDS IN 2D - EQUATIONS OF MOTION

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The equations of motion for ϕ and $\tilde{\phi}$ result to be respectively:

$$\begin{aligned}\partial_\sigma \left[\partial_\sigma \phi - \partial_\tau \tilde{\phi} \right] &= 0 \quad ; \quad \partial_\sigma \left[\partial_\sigma \tilde{\phi} - \partial_\tau \phi \right] = 0 \\ \partial_\sigma \phi - \partial_\tau \tilde{\phi} &= f(\tau) \quad ; \quad \partial_\sigma \tilde{\phi} - \partial_\tau \phi = \tilde{f}(\tau)\end{aligned}\tag{8}$$

FREE SCALAR FIELDS IN 2D - EQUATIONS OF MOTION

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The equations of motion for ϕ and $\tilde{\phi}$ result to be respectively:

$$\begin{aligned}\partial_\sigma \left[\partial_\sigma \phi - \partial_\tau \tilde{\phi} \right] &= 0 \quad ; \quad \partial_\sigma \left[\partial_\sigma \tilde{\phi} - \partial_\tau \phi \right] = 0 \\ \partial_\sigma \phi - \partial_\tau \tilde{\phi} &= f(\tau) \quad ; \quad \partial_\sigma \tilde{\phi} - \partial_\tau \phi = \tilde{f}(\tau)\end{aligned}\quad (8)$$

- Hence, they can be rewritten as first-order equations:

$$\begin{aligned}\partial_\sigma \phi - \partial_\tau \tilde{\phi} &= 0 \\ \partial_\sigma \tilde{\phi} - \partial_\tau \phi &= 0\end{aligned}$$

by invoking another symmetry of \mathcal{L}_{sym} , i.e. the one under a *shift*:

$$\begin{aligned}\phi &\rightarrow \phi + g(\tau) \\ \tilde{\phi} &\rightarrow \tilde{\phi} + \tilde{g}(\tau)\end{aligned}$$

FREE SCALARS FIELDS IN 2D - EQUATIONS OF MOTION

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The equations of motion for ϕ and $\tilde{\phi}$ result to be respectively:

$$\begin{aligned}\partial_\sigma \left[\partial_\sigma \phi - \partial_\tau \tilde{\phi} \right] &= 0 \quad ; \quad \partial_\sigma \left[\partial_\sigma \tilde{\phi} - \partial_\tau \phi \right] = 0 \\ \partial_\sigma \phi - \partial_\tau \tilde{\phi} &= f(\tau) \quad ; \quad \partial_\sigma \tilde{\phi} - \partial_\tau \phi = \tilde{f}(\tau)\end{aligned}\quad (8)$$

- Hence, they can be rewritten as first-order equations:

$$\begin{aligned}\partial_\sigma \phi - \partial_\tau \tilde{\phi} &= 0 \\ \partial_\sigma \tilde{\phi} - \partial_\tau \phi &= 0\end{aligned}$$

by invoking another symmetry of \mathcal{L}_{sym} , i.e. the one under a *shift*:

$$\begin{aligned}\phi &\rightarrow \phi + g(\tau) \\ \tilde{\phi} &\rightarrow \tilde{\phi} + \tilde{g}(\tau)\end{aligned}$$

- The equations of motion reproduce on-shell the duality conditions, after gauging away $f(\tau)$ and $\tilde{f}(\tau)$.

FLOREANINI-JACKIW LAGRANGIANS FOR CHIRAL FIELDS

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The symmetric Lagrangian \mathcal{L}_{sym} can be diagonalized by introducing a pair of scalar fields ϕ_+ and ϕ_- defined by:

$$\phi \equiv \frac{1}{\sqrt{2}} (\phi_+ + \phi_-) \quad ; \quad \tilde{\phi} \equiv \frac{1}{\sqrt{2}} (\phi_+ - \phi_-) \quad (9)$$

in terms of which it becomes the sum of two **Floeanini-Jackiw Lagrangians**, the one associate with ϕ_+ and the other with ϕ_- :

$$\mathcal{L}_{sym} = \mathcal{L}_+(\phi_+) + \mathcal{L}_-(\phi_-) \quad (10)$$

with

$$\mathcal{L}_{\pm}(\phi_{\pm}) = \pm \frac{1}{2} \dot{\phi}_{\pm} \phi'_{\pm} - \frac{1}{2} \phi_{\pm}^{\prime 2} \quad (11)$$

FLOREANINI-JACKIW LAGRANGIANS FOR CHIRAL FIELDS

CLASSICAL AND QUANTUM ASPECTS OF THE STRING DOUBLE SIGMA MODEL

FRANCO PEZZELLA

INTRODUCTION AND MOTIVATION

HODGE-DUAL SYMMETRIC FREE SCALAR FIELDS IN 2D

DOUBLE SIGMA MODEL (CLOSED STRINGS)

DUALITY SYMMETRIC FREE CLOSED STRINGS

QUANTIZATION OF THE DOUBLE STRING MODEL

CONCLUSION AND PERSPECTIVES

- The symmetric Lagrangian \mathcal{L}_{sym} can be diagonalized by introducing a pair of scalar fields ϕ_+ and ϕ_- defined by:

$$\phi \equiv \frac{1}{\sqrt{2}} (\phi_+ + \phi_-) \quad ; \quad \tilde{\phi} \equiv \frac{1}{\sqrt{2}} (\phi_+ - \phi_-) \quad (9)$$

in terms of which it becomes the sum of two **Floeanini-Jackiw Lagrangians**, the one associate with ϕ_+ and the other with ϕ_- :

$$\mathcal{L}_{sym} = \mathcal{L}_+(\phi_+) + \mathcal{L}_-(\phi_-) \quad (10)$$

with

$$\mathcal{L}_\pm(\phi_\pm) = \pm \frac{1}{2} \dot{\phi}_\pm \phi'_\pm - \frac{1}{2} \phi'^2_\pm \quad (11)$$

- It is only on-shell that ϕ_\pm become functions of $\sigma \pm \tau$:

$$\dot{\phi}_+ = \phi'_+ \quad \dot{\phi}_- = -\phi'_- \quad (12)$$

SYMMETRIES

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- \mathcal{L}_{sym} is invariant under **space-time translations** acting as (the constant parameters of the transformations are omitted):

$$\delta_\tau \phi = \dot{\phi} \quad ; \quad \delta_\sigma \phi = \phi' \quad (13)$$

and under **modified global Lorentz transformations**:

$$\delta_L \phi = \tau \phi' + \sigma \tilde{\phi}' \quad ; \quad \delta_L \tilde{\phi} = \tau \tilde{\phi}' + \sigma \phi'$$

that on-shell become the usual two-dimensional Lorentz rotations:

$$\delta_L \phi = \tau \phi' + \sigma \dot{\phi} \quad ; \quad \delta_L \tilde{\phi} = \tau \tilde{\phi}' + \sigma \ddot{\phi}$$

- **The Lorentz invariance is recovered on-shell.**

CHIRAL AND NON-CHIRAL BASIS

- The free Lagrangians considered here can be rewritten, in both cases, as:

$$\mathcal{L}_0 = \frac{1}{2} (C_{ij} \partial_0 \Phi^i \partial_1 \Phi^j + M_{ij} \partial_1 \Phi^i \partial_1 \Phi^j) . \quad (14)$$

In the **chiral basis** $\Phi^i = (\phi_+, \phi_-)$ ($i = 1, 2$)

$$C = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \text{ and } M = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} ;$$

in the **non-chiral basis** $\Phi^i = (\phi, \tilde{\phi})$

$$C \equiv \Omega = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \text{ and } M = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} .$$

CHIRAL AND NON-CHIRAL BASIS

- The free Lagrangians considered here can be rewritten, in both cases, as:

$$\mathcal{L}_0 = \frac{1}{2} (C_{ij} \partial_0 \Phi^i \partial_1 \Phi^j + M_{ij} \partial_1 \Phi^i \partial_1 \Phi^j) . \quad (14)$$

In the **chiral basis** $\Phi^i = (\phi_+, \phi_-)$ ($i = 1, 2$)

$$C = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \text{ and } M = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} ;$$

in the **non-chiral basis** $\Phi^i = (\phi, \tilde{\phi})$

$$C \equiv \Omega = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \text{ and } M = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} .$$

- C and M , will become respectively, in the string case, the $O(D, D)$ invariant metric and the *generalised metric*.

TWO-DIMENSIONAL SCALAR FIELDS ON CURVED SPACE

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- In order to couple \mathcal{L}_{sym} , or the two FJ Lagrangians for chiral scalar fields, to the external 2-bein e^a one is to replace $\partial_a \rightarrow e_a^\alpha$ and to multiply by $e \equiv \det e_\alpha^a$:

$$\begin{aligned} \mathcal{L}_{sym} = & \frac{1}{2} e \left[e_0^\alpha e_1^\beta \partial_n \phi \partial_m \tilde{\phi} + e_1^\alpha e_0^\beta \partial_\alpha \phi \partial_\beta \tilde{\phi} \right. \\ & \left. - e_1^\alpha e_1^\beta \partial_\alpha \phi \partial_\beta \phi - e_1^\alpha e_1^\beta \partial_\alpha \tilde{\phi} \partial_\beta \tilde{\phi} \right] \end{aligned}$$

- After eliminating $\tilde{\phi}$ through its equation of motion, one returns to the usual scalar Lagrangian:

$$\mathcal{L} = \frac{1}{2} e \eta^{ab} e_a^\alpha e_b^\beta \partial_\alpha \phi \partial_\beta \phi \quad (15)$$

GENERAL STRING SIGMA-MODEL

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- General string “sigma model”:

$$S = -\frac{T}{2} \int d^2\sigma e [C_{ij} \nabla_0 \chi^i \nabla_1 \chi^j + M_{ij} \nabla_1 \chi^i \nabla_1 \chi^j] \quad (16)$$

- $e^a_{\alpha} \rightarrow$ zweibein defined on the string world-sheet.

GENERAL STRING SIGMA-MODEL

- General string “sigma model”:

$$S = -\frac{T}{2} \int d^2\sigma e [C_{ij} \nabla_0 \chi^i \nabla_1 \chi^j + M_{ij} \nabla_1 \chi^i \nabla_1 \chi^j] \quad (16)$$

- $e^a{}_\alpha \rightarrow$ zweibein defined on the string world-sheet.
- $C_{ij} = C_{ji}$ and $M_{ij} = M_{ji}$; $\nabla_a \chi^i = e_a^\alpha \partial_\alpha \chi^i$, the functions χ^i the string coordinates in an N -dimensional Riemannian target space.

GENERAL STRING SIGMA-MODEL

- General string “sigma model”:

$$S = -\frac{T}{2} \int d^2\sigma e [C_{ij} \nabla_0 \chi^i \nabla_1 \chi^j + M_{ij} \nabla_1 \chi^i \nabla_1 \chi^j] \quad (16)$$

- $e^a{}_\alpha \rightarrow$ zweibein defined on the string world-sheet.
- $C_{ij} = C_{ji}$ and $M_{ij} = M_{ji}$; $\nabla_a \chi^i = e_a^\alpha \partial_\alpha \chi^i$, the functions χ^i the string coordinates in an N -dimensional Riemannian target space.

Symmetries

- 1 Invariance under **diffeomorphisms**:

$$\sigma^\alpha \rightarrow \sigma'^\alpha(\sigma)$$

- 2 Invariance under **Weyl transformations**:

$$e^a{}_\alpha \rightarrow \lambda(\sigma) e^a{}_\alpha$$

- 3 Request of invariance under **local Lorentz transformations**:

$$e^a{}_\alpha \rightarrow e'^a{}_\alpha = \Lambda^a{}_b(\sigma) e^b{}_\alpha \text{ where } \Lambda^a{}_b \text{ is an arbitrary Lorentz matrix } SO(1, 1).$$

REQUIRING LOCAL LORENTZ INVARIANCE

- The action S is not manifestly invariant under the group $SO(1, 1)$ of local Lorentz transformations:

$$\delta e^a{}_\alpha = \alpha(\sigma)\epsilon^a{}_b(\sigma)e^b{}_\alpha \quad (17)$$

but such invariance has to hold since physical observables are independent on the choice of the vielbein. Hence, **the theory is required to be locally Lorentz invariant on shell.**

REQUIRING LOCAL LORENTZ INVARIANCE

- The action S is not manifestly invariant under the group $SO(1, 1)$ of local Lorentz transformations:

$$\delta e^a{}_\alpha = \alpha(\sigma) \epsilon^a{}_b(\sigma) e^b{}_\alpha \quad (17)$$

but such invariance has to hold since physical observables are independent on the choice of the vielbein. Hence, **the theory is required to be locally Lorentz invariant on shell.**

- Since the variation of S under an infinitesimal local Lorentz transformation results to be:

$$\frac{\delta S}{\delta e^a{}_\alpha} \delta e^a{}_\alpha = \alpha(\sigma) \frac{e}{2} \epsilon^a{}_b t_a{}^b$$

the above requirement implies:

$$\epsilon^{ab} t_{ab} = 0 \quad t_a{}^b \equiv -\frac{2}{T} \frac{1}{e} \frac{\delta S}{\delta e^a{}_\alpha} e^b{}_\alpha \quad (18)$$

- The Weyl invariance implies:

$$\eta^{ab} t_{ab} = 0 .$$

- The Weyl invariance implies:

$$\eta^{ab} t_{ab} = 0 .$$

- The equations of motion for $e^a{}_\alpha$ give $t_{ab} = 0$ providing constraints that have to be imposed at classical and quantum levels, analogously to what happens in the ordinary formulation with $T_{\alpha\beta} = -\frac{2}{T\sqrt{g}} \frac{\delta S}{\delta g^{\alpha\beta}} = 0$. Hence, on the solutions of these equations the local Lorentz invariance holds.

- The Weyl invariance implies:

$$\eta^{ab} t_{ab} = 0 .$$

- The equations of motion for e^a_α give $t_{ab} = 0$ providing constraints that have to be imposed at classical and quantum levels, analogously to what happens in the ordinary formulation with $T_{\alpha\beta} = -\frac{2}{T\sqrt{g}} \frac{\delta S}{\delta g^{\alpha\beta}} = 0$. Hence, on the solutions of these equations the local Lorentz invariance holds.
- Local symmetries (Reparametrization + Weyl + Local Lorentz inv.) allow to fix the **flat gauge**

$$e^a_\alpha = \delta_\alpha^a .$$

CONSTRAINTS AND EQUATIONS OF MOTION

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The constraint $\epsilon^{ab}t_{ab} = 0$ can be rewritten in the following way:

$$\begin{aligned} & [C_{ij}\partial_0\chi^j + M_{ij}\partial_1\chi^j] (C^{-1})^{ik} [C_{kl}\partial_0\chi^l + M_{kl}\partial_1\chi^l] \\ & + [C - MC^{-1}M]_{ij} \partial_1\chi^i \partial_1\chi^j = 0. \end{aligned} \quad (19)$$

CONSTRAINTS AND EQUATIONS OF MOTION

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The constraint $\epsilon^{ab}t_{ab} = 0$ can be rewritten in the following way:

$$\begin{aligned} & [C_{ij}\partial_0\chi^j + M_{ij}\partial_1\chi^j] (C^{-1})^{ik} [C_{kl}\partial_0\chi^l + M_{kl}\partial_1\chi^l] \\ & + [C - MC^{-1}M]_{ij} \partial_1\chi^i \partial_1\chi^j = 0. \end{aligned} \quad (19)$$

- Equations of motion for χ^i :

$$\begin{aligned} & \partial_1 [C_{ij}\partial_0\chi^j + M_{ij}\partial_1\chi^j] - \Gamma^l{}_{ik} C_{lj}\partial_0\chi^j \partial_1\chi^k \\ & - \frac{1}{2}(\partial_i M_{jk})\partial_1\chi^j \partial_1\chi^k = 0 \end{aligned}$$

CONSTRAINTS AND EQUATIONS OF MOTION

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The constraint $\epsilon^{ab}t_{ab} = 0$ can be rewritten in the following way:

$$\begin{aligned} & [C_{ij}\partial_0\chi^j + M_{ij}\partial_1\chi^j] (C^{-1})^{ik} [C_{kl}\partial_0\chi^l + M_{kl}\partial_1\chi^l] \\ & + [C - MC^{-1}M]_{ij} \partial_1\chi^i \partial_1\chi^j = 0. \end{aligned} \quad (19)$$

- Equations of motion for χ^i :

$$\begin{aligned} & \partial_1 [C_{ij}\partial_0\chi^j + M_{ij}\partial_1\chi^j] - \Gamma^l{}_{ik} C_{lj}\partial_0\chi^j \partial_1\chi^k \\ & - \frac{1}{2}(\partial_i M_{jk})\partial_1\chi^j \partial_1\chi^k = 0 \end{aligned}$$

- Boundary conditions:

$$\left[\left(\frac{1}{2} C_{ij}\partial_0\chi^j + M_{ij}\partial_1\chi^j \right) \right]_{\sigma=0}^{\sigma=\pi} = 0$$

CONSTANT BACKGROUNDS

- When C and M are constant, the equations of motion for χ^i drastically simplifies into:

$$\partial_1 [C_{ij}\partial_0\chi^j + M_{ij}\partial_1\chi^j] = 0 .$$

CONSTANT BACKGROUNDS

- When C and M are constant, the equations of motion for χ^i drastically simplifies into:

$$\partial_1 [C_{ij}\partial_0\chi^j + M_{ij}\partial_1\chi^j] = 0 .$$

- The further local gauge invariance of the action under shifts as:

$$\delta\chi^i = f^i(\tau, \sigma) \quad \text{with} \quad \nabla_1 f^i = 0 \quad (20)$$

allows to rewrite the equation of motion for χ^i as:

$$C_{ij}\partial_0\chi^j + M_{ij}\partial_1\chi^j = 0 \quad (21)$$

with boundary conditions dictated by the vanishing of the surface integral:

$$\frac{1}{2} \int d\tau C_{ij} [\partial_0\chi^j \delta\chi^i] \Big|_{\sigma=0}^{\sigma=\tau} = 0 \quad (22)$$

describing both open strings with Dirichlet boundary conditions and closed strings.

EMERGING OUT OF $O(D, D)$

- This causes the constraint on the ϵ -trace to become:

$$[C - MC^{-1}M]_{ij} \partial_1 \chi^i \partial_1 \chi^j = 0$$

implying the restriction on C and M : $C = MCM$.

- After rotating and rescaling χ^i , C can always be put in the diagonal form:

$$C = (1, \dots, 1, -1, \dots, -1)$$

with N_+ eigenvalues 1 and N_- eigenvalues -1 and $N = N_+ + N_-$.

So the action can be interpreted as describing N_+ chiral and N_- antichiral scalars interacting via the bilinear term

$(M_{ij} + \delta_{ij}) \nabla_1 \chi^i \nabla_1 \chi^j$ and the absence of a quantum Lorentz anomaly requires $N_+ = N_- = D = \frac{N}{2}$. Hence, $N = 2D$.

- C becomes the $O(D, D)$ invariant metric while $C = MCM$ implies that M is an $O(D, D)$ element.

NON-CHIRAL COORDINATES

- It is possible to make a change of coordinates in the $2D$ -dimensional target space according to the definition:

$$X^\mu \equiv \frac{1}{\sqrt{2}} (X_+^\mu + X_-^\mu) \quad ; \quad \tilde{X}_\mu \equiv \delta_{\mu\nu} \frac{1}{\sqrt{2}} (X_+^\nu - X_-^\nu)$$

NON-CHIRAL COORDINATES

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- It is possible to make a change of coordinates in the $2D$ -dimensional target space according to the definition:

$$X^\mu \equiv \frac{1}{\sqrt{2}} (X_+^\mu + X_-^\mu) \quad ; \quad \tilde{X}_\mu \equiv \delta_{\mu\nu} \frac{1}{\sqrt{2}} (X_+^\nu - X_-^\nu)$$

- It makes the matrix C become off-diagonal:

$$C_{ij} = -\Omega_{ij} \quad ; \quad \Omega_{ij} = \begin{pmatrix} 0_{\mu\nu} & \mathbb{I}_\mu^\nu \\ \mathbb{I}_\nu^\mu & 0^{\mu\nu} \end{pmatrix}$$

with $(\Omega)_{ij} = (\Omega^{-1})^{ij}$.

NON-CHIRAL COORDINATES

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- It is possible to make a change of coordinates in the $2D$ -dimensional target space according to the definition:

$$X^\mu \equiv \frac{1}{\sqrt{2}} (X_+^\mu + X_-^\mu) \quad ; \quad \tilde{X}_\mu \equiv \delta_{\mu\nu} \frac{1}{\sqrt{2}} (X_+^\nu - X_-^\nu)$$

- It makes the matrix C become off-diagonal:

$$C_{ij} = -\Omega_{ij} \quad ; \quad \Omega_{ij} = \begin{pmatrix} 0_{\mu\nu} & \mathbb{I}_\mu^\nu \\ \mathbb{I}_\nu^\mu & 0^{\mu\nu} \end{pmatrix}$$

with $(\Omega)_{ij} = (\Omega^{-1})^{ij}$.

- The expression for M results to be:

$$M_{ij} = \begin{pmatrix} (G - B G^{-1} B)_{\mu\nu} & (B G^{-1})_\mu^\nu \\ (-G^{-1} B)^\mu_\nu & (G^{-1})^{\mu\nu} \end{pmatrix}$$

being M parametrized by D^2 .

$O(D, D)$ INVARIANCE

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The sigma-model action can be expressed, in the non-chiral basis, as:

$$S = -\frac{T}{2} \int d^2\sigma [\Omega_{ij} \partial_0 \chi^i \partial_1 \chi^j - M_{ij} \partial_1 \chi^i \partial_1 \chi^j].$$

- It is invariant under the combined $O(D, D)$ transformations of χ^i and the matrix of the couplings parameters in M :

$$\chi' = \mathcal{R}\chi ; \quad M' = \mathcal{R}^{-t} M \mathcal{R}^{-1} ; \quad \mathcal{R}^t \Omega \mathcal{R} = \Omega ; \quad \mathcal{R} \in O(D, D).$$

- The $O(D, D)$ invariant metric Ω is itself an element of $O(D, D)$.

RECOVERING THE FAMILIAR T-DUALITY INVARIANCE

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- Define the **duality transformation** $\mathcal{R} = \Omega$ under which $X^\mu \rightarrow \tilde{X}_\mu$. The action, expressed in terms of X^μ and \tilde{X}_μ , after this transformation, becomes:

$$S = -\frac{T}{2} \int d^2\sigma \left[\partial_0 X^\mu \partial_1 \tilde{X}_\mu + \partial_0 \tilde{X}^\mu \partial_1 X_\mu - (G - B G^{-1} B)_{\mu\nu} \partial_1 X^\mu \partial_1 X^\nu - (B G^{-1})_\mu{}^\nu \partial_1 X^\mu \partial_1 \tilde{X}_\nu + (G^{-1} B)^\mu{}_\nu \partial_1 \tilde{X}_\mu \partial_1 X^\nu - (G^{-1})^{\mu\nu} \partial_1 \tilde{X}_\mu \partial_1 \tilde{X}_\nu \right]$$

and exhibits what in string theory is the familiar **T-duality invariance**, in presence of backgrounds, i.e. $X \leftrightarrow \tilde{X}$ together with a transformation of the generalised metric given by $M' = M^{-1}$, i.e.

$$G \leftrightarrow (G - B G^{-1} B)^{-1} \\ B G^{-1} \leftrightarrow -G^{-1} B$$

CORRESPONDENCE WITH THE STANDARD FORMULATION IN CONSTANT BACKGROUNDS

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- In order to understand the relation to the standard formulation, one can integrate over \tilde{X}_μ by eliminating it through the use of the equations of motion. In the case of G, B constant one gets the standard sigma-model action:

$$S[X] = -\frac{T}{2} \int d^2\sigma (\sqrt{G} G^{mm} + \epsilon^{mn})(G + B)_{\mu\nu} \partial_m X^\mu \partial_n X^\nu$$

which describes the toroidal compactification under proper periodicity conditions on X . If, instead, one eliminates X from its equation of motion one obtains the dual model for \tilde{X} :

$$S[\tilde{X}] = -\frac{T}{2} \int d^2\sigma (\sqrt{G} G^{mn} + \epsilon^{mn})(G + B)^{-1\mu\nu} \partial_m \tilde{X}^\mu \partial_n \tilde{X}^\nu$$

- The action $S[X, \tilde{X}]$ is therefore a first-order action which interpolates between $S[X]$ and $S[\tilde{X}]$ and is manifestly duality symmetric.

FREE CLOSED DOUBLED STRINGS

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- From the above formulation it is easy to derive the free action for doubled strings. This corresponds to the case in which:

$$C = - \begin{pmatrix} 0 & \mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix} \quad \text{and} \quad M = \begin{pmatrix} G & 0 \\ 0 & G^{-1} \end{pmatrix} \quad (23)$$

with $G_{\mu\nu}$ being the **flat metric** in the target space. One gets:

$$\begin{aligned} S_0 &= S[X^\mu, e] + S[\tilde{X}_\mu, e] \\ &= -\frac{1}{4\pi\alpha'} \int d^2\sigma e \left[\nabla_0 X^\mu \nabla_1 \tilde{X}_\mu + \nabla_0 \tilde{X}^\mu \nabla_1 X_\mu \right. \\ &\quad \left. - G_{\mu\nu} \nabla_1 X^\mu \nabla_1 X^\nu - \tilde{G}^{\mu\nu} \nabla_1 \tilde{X}_\mu \nabla_1 \tilde{X}_\nu \right] \\ &= S[X_+^\mu, e] + S[X_-^\mu, e] \end{aligned} \quad (24)$$

with $\tilde{G}^{\mu\nu} = G^{-1\mu\nu}$, $\nabla_a = e_a^\alpha \partial_\alpha$ and $\mu = 1, \dots, D$. This is invariant under $X^\mu \leftrightarrow \tilde{X}_\mu$ together with $G_{\mu\nu} \leftrightarrow \tilde{G}^{\mu\nu}$.

INSERTING VERTEX OPERATORS

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The free action S_0 still describes D and not $2D$ scalar degrees of freedom (only the zero mode of X and \tilde{X} are independent on-shell).

INSERTING VERTEX OPERATORS

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The free action S_0 still describes D and not $2D$ scalar degrees of freedom (only the zero mode of X and \tilde{X} are independent on-shell).
- S_0 can be perturbed by $S_{int}[X, \tilde{X}]$ with the insertion of vertex operators involving both X and \tilde{X} . If S_{int} does not depend on \tilde{X} one can integrate \tilde{X} out in the path integral of the theory and reproduce the usual results of the standard formulation.

INSERTING VERTEX OPERATORS

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The free action S_0 still describes D and not $2D$ scalar degrees of freedom (only the zero mode of X and \tilde{X} are independent on-shell).
- S_0 can be perturbed by $S_{int}[X, \tilde{X}]$ with the insertion of vertex operators involving both X and \tilde{X} . If S_{int} does not depend on \tilde{X} one can integrate \tilde{X} out in the path integral of the theory and reproduce the usual results of the standard formulation.
- Assuming that strings are compactified on a circle of radius R , one should expect that: **at large scales** $R \gg \sqrt{\alpha'}$ the relevant interactions are $S_{int}(X)$; **at intermediate scales** $R \equiv \sqrt{\alpha'}$ the relevant interactions involve both X and \tilde{X} while **at small scales** $R \ll \sqrt{\alpha'}$ the relevant interactions are $S_{int}(\tilde{X})$.

INSERTING VERTEX OPERATORS

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The free action S_0 still describes D and not $2D$ scalar degrees of freedom (only the zero mode of X and \tilde{X} are independent on-shell).
- S_0 can be perturbed by $S_{int}[X, \tilde{X}]$ with the insertion of vertex operators involving both X and \tilde{X} . If S_{int} does not depend on \tilde{X} one can integrate \tilde{X} out in the path integral of the theory and reproduce the usual results of the standard formulation.
- Assuming that strings are compactified on a circle of radius R , one should expect that: **at large scales** $R \gg \sqrt{\alpha'}$ the relevant interactions are $S_{int}(X)$; **at intermediate scales** $R \equiv \sqrt{\alpha'}$ the relevant interactions involve both X and \tilde{X} while at **$R \ll \sqrt{\alpha'}$** the relevant interactions are $S_{int}(\tilde{X})$.
- **The duality symmetric formulation may be considered as a natural generalization of the standard one at the string scale.**

EQUIVALENCE BETWEEN NON-COVARIANT AND COVARIANT ACTIONS

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The action

$$S = -\frac{T}{2} \int d^2\sigma [C_{ij}\partial_0\chi^i \partial_1\chi^j + M_{ij}\partial_1\chi^i \partial_1\chi^j] \quad (25)$$

is candidate to describe, with C and M constant, bosonic closed strings in a background and compactified on a torus T^D . It exhibits a manifest T -duality invariance $O(D, D)$ with the fields χ^i interpreted as string coordinates on the double torus T^{2D} .

EQUIVALENCE BETWEEN NON-COVARIANT AND COVARIANT ACTIONS

CLASSICAL AND QUANTUM ASPECTS OF THE STRING DOUBLE SIGMA MODEL

FRANCO PEZZELLA

INTRODUCTION AND MOTIVATION

HODGE-DUAL SYMMETRIC FREE SCALAR FIELDS IN 2D

DOUBLE SIGMA MODEL (CLOSED STRINGS)

DUALITY SYMMETRIC FREE CLOSED STRINGS

QUANTIZATION OF THE DOUBLE STRING MODEL

CONCLUSION AND PERSPECTIVES

- The action

$$S = -\frac{T}{2} \int d^2\sigma [C_{ij}\partial_0\chi^i\partial_1\chi^j + M_{ij}\partial_1\chi^i\partial_1\chi^j] \quad (25)$$

is candidate to describe, with C and M constant, bosonic closed strings in a background and compactified on a torus T^D . It exhibits a manifest T -duality invariance $O(D, D)$ with the fields χ^i interpreted as string coordinates on the double torus T^{2D} .

- It can be shown to be equivalent to the following **covariant action** (Hull, 2005):

$$S = -\frac{T}{2} \int d^2\sigma \partial^\alpha\chi^i M_{ij} \partial_\alpha\chi^j \quad (26)$$

with the self-duality relation imposed in order to halve the degrees of freedom from $2D$ to D (also Duff, 1987):

$$\partial_\alpha\chi^i = \epsilon_{\alpha\beta}\eta^{ij}M_{jk}(\partial^\beta\chi^k) \quad (27)$$

equivalent to the condition $\epsilon_{ab}t^{ab} = 0$.

EQUIVALENCE BETWEEN NON-COVARIANT AND COVARIANT ACTIONS

CLASSICAL AND QUANTUM ASPECTS OF THE STRING DOUBLE SIGMA MODEL

FRANCO PEZZELLA

INTRODUCTION AND MOTIVATION

HODGE-DUAL SYMMETRIC FREE SCALAR FIELDS IN 2D

DOUBLE SIGMA MODEL (CLOSED STRINGS)

DUALITY SYMMETRIC FREE CLOSED STRINGS

QUANTIZATION OF THE DOUBLE STRING MODEL

CONCLUSION AND PERSPECTIVES

- The action

$$S = -\frac{T}{2} \int d^2\sigma [C_{ij}\partial_0\chi^i\partial_1\chi^j + M_{ij}\partial_1\chi^i\partial_1\chi^j] \quad (25)$$

is candidate to describe, with C and M constant, bosonic closed strings in a background and compactified on a torus T^D . It exhibits a manifest T -duality invariance $O(D, D)$ with the fields χ^i interpreted as string coordinates on the double torus T^{2D} .

- It can be shown to be equivalent to the following **covariant action** (Hull, 2005):

$$S = -\frac{T}{2} \int d^2\sigma \partial^\alpha\chi^i M_{ij} \partial_\alpha\chi^j \quad (26)$$

with the self-duality relation imposed in order to halve the degrees of freedom from $2D$ to D (also Duff, 1987):

$$\partial_\alpha\chi^i = \epsilon_{\alpha\beta}\eta^{ij}M_{jk}(\partial^\beta\chi^k) \quad (27)$$

equivalent to the condition $\epsilon_{ab}t^{ab} = 0$.

NON-CONSTANT BACKGROUNDS

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- Aim: to introduce interactions and understand if the local Lorentz constraint still holds under the form $C = MCM$ in case of non-constant backgrounds.

FIRST CASE:

C constant and M only X -dependent (or only \bar{X} -dependent).

- In the case in which $C = \Omega$ and M only X -dependent, in deriving the equations of motion for X^μ and \tilde{X}_μ one has to keep in consideration the contribution coming from the term

$$\frac{1}{2}(\partial_i M_{jk})\partial_1 \chi^j \partial_1 \chi^k.$$

- The equations of motion for X^μ and \tilde{X}_μ respectively become:

$$\partial_1 \left[-\partial_0 \tilde{X}_\mu + (G - BG^{-1}B)_{\mu\nu} \partial_1 X^\nu + (BG^{-1})_\mu^\nu \partial_1 \tilde{X}_\nu \right] = \frac{1}{2} \partial_1 X^\nu \left[\partial_\mu (G - BG^{-1}B)_{\nu\rho} \partial_1 X^\rho + \partial_\mu (BG^{-1})^{\nu\rho} \partial_1 \tilde{X}_\rho \right]$$

and

$$\partial_1 \left[-\partial_0 X^\mu + (-G^{-1}B)^\mu_\nu \partial_1 X^\nu + (G^{-1})^{\mu\nu} \partial_1 \tilde{X}_\nu \right] = \frac{1}{2} \partial_1 \tilde{X}_\nu \left[\bar{\partial}^\mu (-G^{-1}B)^\nu_\rho \partial_1 X^\rho + \bar{\partial}^\mu (G^{-1})^{\nu\rho} \partial_1 \tilde{X}_\rho \right] = 0$$

where $\bar{\partial}^\mu$ denotes the derivative with respect to \tilde{X}_μ .

- Also in this case, one can use the invariance of the equation of motion for \tilde{X}_μ under *shifts* for putting:

$$-\partial_0 X^\mu + (-G^{-1}B)^\mu{}_\nu \partial_1 X^\nu + (G^{-1})^{\mu\nu} \partial_1 \tilde{X}_\nu = 0. \quad (28)$$

- Also in this case, one can use the invariance of the equation of motion for \tilde{X}_μ under *shifts* for putting:

$$-\partial_0 X^\mu + (-G^{-1}B)^\mu{}_\nu \partial_1 X^\nu + (G^{-1})^{\mu\nu} \partial_1 \tilde{X}_\nu = 0. \quad (28)$$

- When this expression is substituted in the condition $\epsilon_{ab} t^{ab} = 0$, that is valid for any kind of backgrounds:

$$\begin{aligned} [C_{ij} \partial_0 \chi^j + M_{ij} \partial_1 \chi^j] (C^{-1})^{ik} [C_{kl} \partial_0 \chi^l + M_{kl} \partial_1 \chi^l] \\ + [C - MC^{-1}M]_{ij} \partial_1 \chi^i \partial_1 \chi^j = 0. \end{aligned} \quad (29)$$

one can easily see that the off-diagonal structure of C makes the first term vanish and so one gets again the condition $C = MCM$ characterizing the $O(D, D)$ invariance.

- Also in this case, one can use the invariance of the equation of motion for \tilde{X}_μ under *shifts* for putting:

$$-\partial_0 X^\mu + (-G^{-1}B)^\mu{}_\nu \partial_1 X^\nu + (G^{-1})^{\mu\nu} \partial_1 \tilde{X}_\nu = 0. \quad (28)$$

- When this expression is substituted in the condition $\epsilon_{ab} t^{ab} = 0$, that is valid for any kind of backgrounds:

$$\begin{aligned} [C_{ij} \partial_0 \chi^j + M_{ij} \partial_1 \chi^j] (C^{-1})^{ik} [C_{kl} \partial_0 \chi^l + M_{kl} \partial_1 \chi^l] \\ + [C - MC^{-1}M]_{ij} \partial_1 \chi^i \partial_1 \chi^j = 0. \end{aligned} \quad (29)$$

one can easily see that the off-diagonal structure of C makes the first term vanish and so one gets again the condition $C = MCM$ characterizing the $O(D, D)$ invariance.

- The same result is obtained if one considers $C = \Omega$ and M only \tilde{X} -dependent.

- Also in this case, one can use the invariance of the equation of motion for \tilde{X}_μ under *shifts* for putting:

$$-\partial_0 X^\mu + (-G^{-1}B)^\mu_\nu \partial_1 X^\nu + (G^{-1})^{\mu\nu} \partial_1 \tilde{X}_\nu = 0. \quad (28)$$

- When this expression is substituted in the condition $\epsilon_{ab} t^{ab} = 0$, that is valid for any kind of backgrounds:

$$\begin{aligned} [C_{ij} \partial_0 \chi^j + M_{ij} \partial_1 \chi^j] (C^{-1})^{ik} [C_{kl} \partial_0 \chi^l + M_{kl} \partial_1 \chi^l] \\ + [C - MC^{-1}M]_{ij} \partial_1 \chi^i \partial_1 \chi^j = 0. \end{aligned} \quad (29)$$

one can easily see that the off-diagonal structure of C makes the first term vanish and so one gets again the condition $C = MCM$ characterizing the $O(D, D)$ invariance.

- The same result is obtained if one considers $C = \Omega$ and M only \tilde{X} -dependent.

FURTHER OSSERVATION ON $C = \Omega$ AND $M = M(X)$

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- In the case of $C = \Omega$, $M = M(X)$ the constraint $C = MCM$ is still valid and the expression for M keeps on being:

$$M_{ij} = \begin{pmatrix} (G - B G^{-1} B)_{\mu\nu} & (B G^{-1})_{\mu}^{\nu} \\ (-G^{-1} B)^{\mu}_{\nu} & (G^{-1})^{\mu\nu} \end{pmatrix}$$

but now with X -dependent G and B .

- Starting from $S(X^{\mu}, \tilde{X}_{\mu})$ and eliminating \tilde{X}_{μ} through the equation of motion, one can get the usual sigma-model action for X^{μ} :

$$S[X] = -\frac{T}{2} \int d^2\sigma (\sqrt{g} g^{ab} + \epsilon^{ab}) (G + B)_{\mu\nu} \partial_a X^{\mu} \partial_b X^{\nu} \quad (30)$$

that corresponds to the usual formulation of the world sheet of the string in arbitrary background $(G + B)$.

- If X^{μ} is eliminated, then one gets the dual sigma model for \tilde{X}_{μ} :

$$S[\tilde{X}] = -\frac{T}{2} \int d^2\sigma (\sqrt{g} g^{ab} + \epsilon^{ab}) (G + B)^{-1\mu\nu} \partial_a \tilde{X}_{\mu} \partial_b \tilde{X}_{\nu} \quad (31)$$

C AND M BOTH NON-CONSTANT

SECOND CASE:

C and M both depending only on X (or \bar{X}).

- In this case one has to consider, in the equation of motion for \tilde{X}_μ , also the contribution coming from

$$-\Gamma^l{}_{ik} C_{ij} \partial_0 \chi^j \partial_1 \chi^k$$

C AND M BOTH NON-CONSTANT

SECOND CASE:

C and M both depending only on X (or \bar{X}).

- In this case one has to consider, in the equation of motion for \tilde{X}_μ , also the contribution coming from

$$-\Gamma^l{}_{ik} C_{lj} \partial_0 \chi^j \partial_1 \bar{\chi}^k$$

- When rewritten explicitly, this quantity vanishes when the index runs over the ones of \tilde{X}_μ and therefore it does not give any contribution to the equation of motion of this coordinate.

C AND M BOTH NON-CONSTANT

SECOND CASE:

C and M both depending only on X (or \bar{X}).

- In this case one has to consider, in the equation of motion for \tilde{X}_μ , also the contribution coming from

$$-\Gamma^l{}_{ik} C_{ij} \partial_0 \chi^j \partial_1 \chi^k$$

- When rewritten explicitly, this quantity vanishes when the index runs over the ones of \tilde{X}_μ and therefore it does not give any contribution to the equation of motion of this coordinate.
- One can conclude that the condition $C = MCM$ still holds under the hypothesis that C and/or M are dependent only on X (or only on \tilde{X}).

C AND M BOTH X, \tilde{X} -DEPENDENT

THIRD CASE:

both C and M depending on the coordinates χ^i .

- One can think to introduce a small parameter $\epsilon = \frac{\sqrt{\alpha'}}{r_c}$ and to expand C and M up to the second order according to:

$$\begin{aligned}C &= C_0 + \epsilon C_1 + \epsilon^2 C_2 \\M &= M_0 + \epsilon M_1 + \epsilon^2 M_2\end{aligned}\tag{32}$$

- By linearizing the condition $\epsilon_{ab} t^{ab} = 0$ and the equations of motion for the coordinates, one gets, **at the order ϵ** :

$$(\epsilon_{ab} t^{ab})_{\text{on-shell}} = -\frac{1}{2} Q_{ij} \partial_1 \chi^i \partial_1 \chi^j + O(\epsilon)$$

$$\begin{aligned}Q &= C_1 - (C_0^{-1} M_0)^t M_1 - M_1 (C_0^{-1} M_0) \\&\quad + (C_0^{-1} M_0)^t C_1 (C_0^{-1} M_0)\end{aligned}$$

Hence, the linearized condition on C_0 and M_0 is $Q = 0$.

This condition can be actually derived by linearizing the condition $C = MCM$. So at this order the $O(D, D)$ condition keeps on holding, being the first term in the expression of the ϵ -trace order ϵ^2 :

$$\begin{aligned} [C_{ij}\partial_0\chi^j + M_{ij}\partial_1\chi^j] (C^{-1})^{ik} [C_{kl}\partial_0\chi^l + M_{kl}\partial_1\chi^l] \\ + [C - MC^{-1}M]_{ij} \partial_1\chi^i \partial_1\chi^j = 0. \end{aligned} \quad (33)$$

This means that the latter plays a role going to the order ϵ^2 and the contribution coming from it adds to the one coming from the term proportional to $C - MCM$. Starting from this order, it seems that the $O(D, D)$ invariance does not hold anymore or one can ask if the deformation is compatible with $O(D, D)$ (discussions with Olaf Hohm and Hai Lin). This is the case that seems to reproduce the α' -corrections found in double field theory (Hohm and Zwiebach, 2014)

CONSTRAINTS OF THE FJ LAGRANGIAN

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The quantization of the action in the flat gauge and for constant backgrounds corresponds to the quantization of the Floreanini-Jackiw Lagrangians.

CONSTRAINTS OF THE FJ LAGRANGIAN

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The quantization of the action in the flat gauge and for constant backgrounds corresponds to the quantization of the Floreanini-Jackiw Lagrangians.
- In the case of a discrete number of degrees of freedom q^i with $i = 1, \dots, N$ it looks like:

$$L = \frac{1}{2} q^i c_{ij} \dot{q}^j - V(q) \text{ with } \det c_{ij} \neq 0.$$

CONSTRAINTS OF THE FJ LAGRANGIAN

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- The quantization of the action in the flat gauge and for constant backgrounds corresponds to the quantization of the Floreanini-Jackiw Lagrangians.
- In the case of a discrete number of degrees of freedom q^i with $i = 1, \dots, N$ it looks like:

$$L = \frac{1}{2} q^i c_{ij} \dot{q}^j - V(q) \text{ with } \det c_{ij} \neq 0.$$

- It is first-order and is characterized by N primary second-class constraints:

$$T_j \equiv p_j - \frac{1}{2} q^i c_{ij} \quad (34)$$

with

$$\{T_i, T_j\} = c_{ij} \neq 0$$

QUANTIZATION OF THE FJ LAGRANGIANS

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- In order to quantize the theory, the Dirac quantization method has to be applied with the corresponding brackets:

$$\{f, g\}_{DB} \equiv \{f, T_j\}_{DB} c^{(-1)jk} \{T_k, g\}_{PB}$$

QUANTIZATION OF THE FJ LAGRANGIANS

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- In order to quantize the theory, the Dirac quantization method has to be applied with the corresponding brackets:

$$\{f, g\}_{DB} \equiv \{f, T_j\}_{DB} c^{(-1)jk} \{T_k, g\}_{PB}$$

- According to the usual transition rule $i\{f, g\}_{DB} \rightarrow \{f, g\}$ from the classical to the quantum theory, the following commutators are obtained:

$$[q_i, q_j] = ic_{ij}^{-1} ; [q_i, p_j] = \frac{1}{2}i\delta_{ij} ; [p_i, p_j] = -\frac{1}{4}ic_{ij}$$

QUANTIZATION OF THE DOUBLE SIGMA MODEL - NON COMMUTATIVITY

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- When translated to the string case, one gets, among the others, a non-commutativity relation between X^μ and \tilde{X}_μ :

$$\left[X(\tau, \sigma), \tilde{X}(\tau, \sigma') \right] = \frac{i}{T} \mathbb{I} \epsilon(\sigma - \sigma') \quad (35)$$

with $\epsilon(\sigma) \equiv \frac{1}{2} [\theta(\sigma) - \theta(-\sigma)]$.

- The Dirac quantization method implies that X^μ and \tilde{X}_μ behave like **non-commuting phase space type coordinates**, but it can be shown that their expressions in terms of Fourier modes generate the usual oscillator algebra of the standard formulation (De Angelis, Gionti, Marotta, FP - 2014).

QUANTIZATION OF THE DOUBLE SIGMA MODEL - NON COMMUTATIVITY

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- When translated to the string case, one gets, among the others, a non-commutativity relation between X^μ and \tilde{X}_μ :

$$\left[X(\tau, \sigma), \tilde{X}(\tau, \sigma') \right] = \frac{i}{T} \mathbb{I} \epsilon(\sigma - \sigma') \quad (35)$$

with $\epsilon(\sigma) \equiv \frac{1}{2} [\theta(\sigma) - \theta(-\sigma)]$.

- The Dirac quantization method implies that X^μ and \tilde{X}_μ behave like **non-commuting phase space type coordinates**, but it can be shown that their expressions in terms of Fourier modes generate the usual oscillator algebra of the standard formulation (De Angelis, Gionti, Marotta, FP - 2014).
- From this perspective, this non-commutativity may lead to the interpretation of high-energy scattering in the X -space as effectively "probing" the \tilde{X} -space.

QUANTIZATION OF THE DOUBLE SIGMA MODEL - NON COMMUTATIVITY

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- When translated to the string case, one gets, among the others, a non-commutativity relation between X^μ and \tilde{X}_μ :

$$\left[X(\tau, \sigma), \tilde{X}(\tau, \sigma') \right] = \frac{i}{T} \mathbb{I} \epsilon(\sigma - \sigma') \quad (35)$$

with $\epsilon(\sigma) \equiv \frac{1}{2} [\theta(\sigma) - \theta(-\sigma)]$.

- The Dirac quantization method implies that X^μ and \tilde{X}_μ behave like **non-commuting phase space type coordinates**, but it can be shown that their expressions in terms of Fourier modes generate the usual oscillator algebra of the standard formulation (De Angelis, Gionti, Marotta, FP - 2014).
- From this perspective, this non-commutativity may lead to the interpretation of high-energy scattering in the X -space as effectively "probing" the \tilde{X} -space.

CONCLUSION

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- An $O(D, D)$ manifest formulation has been analyzed, providing a generalization of the standard formulation at the string scale. It is based on the Floreanini-Jackiw Lagrangians for chiral and antichiral scalar fields.

CONCLUSION

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- An $O(D, D)$ manifest formulation has been analyzed, providing a generalization of the standard formulation at the string scale. It is based on the Floreanini-Jackiw Lagrangians for chiral and antichiral scalar fields.
- The $O(D, D; \mathbb{Z})$ T-duality invariance naturally emerges out in the case of toroidal compactifications.

CONCLUSION

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- An $O(D, D)$ manifest formulation has been analyzed, providing a generalization of the standard formulation at the string scale. It is based on the Floreanini-Jackiw Lagrangians for chiral and antichiral scalar fields.
- The $O(D, D; \mathbb{Z})$ T-duality invariance naturally emerges out in the case of toroidal compactifications.
- A doubling of the string coordinates is naturally required and the quantization requires a non-commuting geometry.

CONCLUSION

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- An $O(D, D)$ manifest formulation has been analyzed, providing a generalization of the standard formulation at the string scale. It is based on the Floreanini-Jackiw Lagrangians for chiral and antichiral scalar fields.
- The $O(D, D; \mathbb{Z})$ T-duality invariance naturally emerges out in the case of toroidal compactifications.
- A doubling of the string coordinates is naturally required and the quantization requires a non-commuting geometry.

- Vertex Operators and Scattering Amplitudes

- Vertex Operators and Scattering Amplitudes
- Effective Action through Beta Functions and relation with DFT

PERSPECTIVES

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

- Vertex Operators and Scattering Amplitudes
- Effective Action through Beta Functions and relation with DFT
- Supersymmetric extension

- Vertex Operators and Scattering Amplitudes
- Effective Action through Beta Functions and relation with DFT
- Supersymmetric extension
- Study of the underlying Generalised Geometry

- Vertex Operators and Scattering Amplitudes
- Effective Action through Beta Functions and relation with DFT
- Supersymmetric extension
- Study of the underlying Generalised Geometry

THE END

CLASSICAL AND
QUANTUM
ASPECTS OF THE
STRING DOUBLE
SIGMA MODEL

FRANCO
PEZZELLA

INTRODUCTION
AND MOTIVATION

HODGE-DUAL
SYMMETRIC
FREE SCALAR
FIELDS IN 2D

DOUBLE SIGMA
MODEL (CLOSED
STRINGS)

DUALITY
SYMMETRIC
FREE CLOSED
STRINGS

QUANTIZATION
OF THE DOUBLE
STRING MODEL

CONCLUSION AND
PERSPECTIVES

Thank you for your attention.