

The importance of quantum loops for astrophysical ALPs

Eike Müller October 2022 @ Youngstars

Based on

- Do Direct Detection Experiments Constrain Axionlike Particles Coupled to Electrons?, Ricardo Z. Ferreira, M. C. David Marsh, and EM, Phys. Rev. Lett. 128, 221302
- Strong supernovae bounds on ALPs from quantum loops, Ricardo Z. Ferreira, M.C. David Marsh, and EM, arXiv:2205.07896 (submitted to JCAP)

Introduction: Axionlike particles



- AllPs are naturally light, weakly interacting pseudoscalar particles that appear in many BSM theories
- Att low energies , Eatt the strugge is a series of the they strugge is a series of the series of the
- Inthisstalk: study just two parameters of the EFT
 phenomenologically at the one-loop level (no model building)

$$\mathcal{L}_{\rm EFT} \supset -\frac{1}{2}a(\Box + m_a^2)a + \hat{g}_{ae}(\partial_{\mu}a)\,\bar{\psi}_e\gamma^{\mu}\gamma_5\psi_e + \frac{g_{a}}{4}aFF$$

Outline



Theoretical basis:

1. Effective, one-loop ALP-photon coupling

Phenomenlogical applications:

- 2. Instability of heavy ALP dark matter
- 3. Supernova bounds at one loop



Theoretical basis

Effective, one-loop ALP-photon coupling

The effective ALP-photon coupling



The one-loop, off-shell matrix element has the same structure as the tree-level version:



The effective ALP-photon coupling



Known for a while: the effective coupling *on-shell*, i.e. in a decay process

$$g_{a\gamma}^{(\mathrm{D})} \equiv g_{a\gamma}^{\mathrm{eff}}(q_1^2 = q_2^2 = 0, p^2 = m_a^2) = \frac{2\alpha}{\pi} \hat{g}_{ae} \left[1 - \frac{4m_e^2}{m_a^2} f^2 \left(\frac{4m_e^2}{m_a^2} \right) \right]$$

$$= -\frac{\alpha \hat{g}_{ae}}{\sqrt{6\pi}} \left(\frac{m_a}{m_a} \right)^2 + \mathcal{O} \left(\frac{m_a}{m_a} \right)^4$$
This effective coupling vanishes for massless ALPs, but it is only the right for results of the right for the right based on the right ba

If a photon in the t-channel is off-shell, we get the effective Primakoff coupling:

$$g_{a\gamma}^{(P)} \equiv g_{a\gamma}^{\text{eff}}(q_1^2 = 0, q_2^2 = t, p^2 = m_a^2) = \frac{2\alpha}{\pi} \hat{g}_{ae} \left\{ 1 + \frac{4m_e^2}{m_a^2 - t} \left[f^2 \left(\frac{4m_e^2}{t} \right) - f^2 \left(\frac{4m_e^2}{m_a^2} \right) \right] \right\}$$
$$= \frac{2\alpha}{\pi} \hat{g}_{ae} \left[1 + \frac{4m_e^2}{m_a^2 - t} f^2 \left(\frac{4m_e^2}{t} \right) \right] + \mathcal{O} \left(\frac{m_a}{m_e} \right)^2$$



The effective ALP-photon coupling

The effective coupling can be used in all processes involving ALPs. Further phenomenologically relevant examples include:

ALP-strahlung

 $g_{a\gamma}^{\text{eff}}(q_1^2 = (k_1 - k_2)^2, q_2^2 = (k_3 - k_4)^2, p^2 = m_a^2)$

Phenomenlogical applications

Instability of heavy ALP dark matter

Instability of ALP DM

Asimple consequence: also ALPs only coupled to electrons with $decay_{m_{a}} < 2m_{e}$ decay (into photons)

$$\begin{array}{ccc} & & & & & & \\ & & & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & &$$

 \rightarrow ALP dark matter in the keV mass range is unstable

Instability of ALP DM

Ferreira, Marsh, **EM**, PRL 128 (2022) 221302

Phenomenlogical applications

Supernova bounds at one loop

Electron Bremsstrahlung

Electron-positron fusion

production

absorption

Supernova bounds at one loop: Cooling bound

Duration of SN1987A's neutrino burst constraints the ALP luminosity:

 $L_{\nu} = 3 \cdot 10^{52} \text{ erg s}^{-1} > L_a = 4\pi \int_0^{\kappa_{\nu}} \mathrm{d}r \, r^2 \int_m^{\infty} \mathrm{d}\omega_a \, \omega_a \, \frac{\mathrm{d}^2 n_a^{\mathrm{tot}}}{\mathrm{d}t \, \mathrm{d}\omega_a} \, e^{-\tau(\omega_a, r)}$ absorptior We use the Agile-Boltztran SN Zemodel from Fischer et al., 14/19PRD 104 (2021) 103012

Supernova bounds at one loop: Decay bound

Supernova bounds at one loop: Decay bound

Total production spectrum

$$\mathrm{d}F_{\gamma} = 2 \cdot \mathrm{BR}_{a \to \gamma}$$

$$\omega \cdot f_{c_{\alpha}}(\omega, c_{\alpha}) \mathrm{d}c_{\alpha}$$

Distribution of decay angles

 $-L/l_a(\omega)$ ALP decay length

Following Jaffe and Turner, PRD 55 (1997) 7951-7959

Constnaints, such as:

 $\Theta_{\rm cons.}(\omega, c_{\alpha}, L)$

- The ALLP should not decay inside the SNN programitor
- One can construct a triangle out of $L, d_{SN}, \cos \alpha$
- ~ The energy of the rara is in the range of the detector
- The ray does not a mive a tetethan 223 safeet ether neutrino burst

Integrate numerically over to get the fluence of trayseofther detector

See also Jaeckel, Malta, Redondo Phys.Rev.D 98 (2018) 5, 055032

Ferreira, Marsh, **EM**, 2205.07896

Supernova bounds at one loop

Ferreira, Marsh, **EM**, 2205.07896

Summary

- · Can define an effective ALP-photon coupling at one-loop
- The coupling depends on the process in which it appears (e.g. decay or Primakoff)
- 40000 induced decays place extremely strong bounds on ALP DM, and even exclude it for large masses/couplings
- Using the effective coupling at one loop, we can place the strongest bounds so far on f_{ae} mrSh1987A

Thanks for your attention!

Back Up

Motivation – Why are loops relevant?

Can the ALP interact much more strongly with electrons than with photons?

Theoretically: quantum loops yield a contribution $g_{a\gamma}^{\text{eff}} \sim 10^{-2} \hat{g}_{ae}$

ALPs from SN1987A: two bounds

The neutrino burst of SN1987A would be shortened by ALPs, unless

$$L_a \lesssim L_{\nu} \simeq 3 \times 10^{52} \frac{\mathrm{erg}}{\mathrm{s}}$$

Gamma rays from decaying ALPs would have been detected near earth after the neutrino burst of SN1987A, unless

ALPs from SN1987A

SN model from: T. Fischer, P. Carenza, B. Fore, M. Giannotti, A. Mirizzi and S. Reddy, Observable signatures of enhanced axion emission from protoneutron stars, Phys. Rev. D 104 (2021) 103012, [2108.13726]

ALPs from SN1987A: Reabsorption

For large couplings, reabsorption of ALPs via inverse processes becomes important

$$L_{a} = \int^{R_{\nu}} d^{3}r \int_{m_{a}}^{\infty} d\omega \, \omega \frac{d\dot{n}}{d\omega} e^{-\int_{r}^{R_{\text{far}}} \Delta(\tilde{r},\omega)} \text{Mean free path of the ALPs}$$
For ,nthermal technique the thresh pathetism free quantities technique technique

Outlook & future work

- One can also derive a bound on the total energy deposited into the progenitor's plasma by ALPS sy way to close the gap between cooling & decay bound
- A similar analysis can be done for ALPs predominantly coupling to muons (this was already done, but only with the effective *decay* coupling)
- There are open questions regarding electron propagation and the ALPelectron interaction in hot and dense plasmas thermal field theory problem
- Use these results as input for SN simulations, including ALPs

ALP-fermion interactions

$$\mathcal{L}_{aQED} = -\frac{1}{2}a(\Box + m_a^2)a - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \bar{\psi}_e(i\not\!\!D - m_e)\psi_e + \frac{g_{a\gamma}}{4}aF_{\mu\nu}\tilde{F}^{\mu\nu} + \hat{g}_{ae}(\partial_{\mu}a)\bar{\psi}_e\gamma^{\mu}\gamma_5\psi_e = -\frac{1}{2}a(\Box + m_a^2)a - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \bar{\psi}'_e(i\not\!\!D - m_e)\psi'_e + \frac{1}{4}\left(g_{a\gamma} + \frac{2\alpha}{\pi}\hat{g}_{ae}\right)aF_{\mu\nu}\tilde{F}^{\mu\nu} - i\underbrace{2m_e\hat{g}_{ae}}_{\equiv g_{ae}}a\,\bar{\psi}'_e\gamma_5\psi'_e + \mathcal{O}(\hat{g}_{ae}^2)$$

 $\psi_e = e^{i\hat{g}_{ae}a\gamma_5}\psi'_e$

ALP-electron interactions in a plasma

• Calculating the bremsstrahlung matrix element with a pseudoscalar ALP-electron interaction yields:

Taking plasma effects into account

 $\equiv 2m_e \hat{g}_{ae} f(m_e^{\text{eff}}, \dots)$ • On the other hand, since the pseudoscalar and derivative

 $\mathcal{M}_{\text{brems}}^{\text{scalar}} = g_{ae} f(m_e^{\text{eff}}, \dots)$

interactions lead (in vacuum) to the same matrix element:

 $\mathcal{M}_{\mathrm{brems}}^{\mathrm{derivative}} = 2m_e^{\mathrm{eff}} \hat{g}_{ae} f(m_e^{\mathrm{eff}}, \dots)$ Therefore, apparently $\mathcal{M}_{\mathrm{brems}}^{\mathrm{derivative}} \neq \mathcal{M}_{\mathrm{brems}}^{\mathrm{splasma.}}$ Why is that?

The effective photon couplings

 $g_{a\gamma}^{\text{eff}}(q_1, q_2)$ depends on the 4-momenta of the photons The effective coupling is different in every physical process!

$rac{\mathbf{m}_{e} \ll \mathbf{m}_{e}}{q_{1}^{2}}$ q_{2}^{2} g_{sr}^{eff}	ALP to photon decay 0 0 $-\frac{a}{6\pi} \left(\frac{m_e}{m_e}\right)^2 \hat{g}_{ax}$	$\begin{array}{l} \label{eq:primakoff effect, with $\omega > m_e$} \\ 0 \\ -2\omega^2(1-\cos\theta) = t \\ \frac{2\alpha}{\pi} \hat{g}_{\alpha e} + \mathcal{O}\left(\frac{m_e^2}{\omega^2}\right) \end{array}$	Α	LP to pho	ton decay	$egin{array}{c} m_a \ll m_c \ q_1^2 \ q_2^2 \ g_{ay}^{eff} \end{array}$	ALP to photon decay 0 $-\frac{\alpha}{6\pi} \left(\frac{m_a}{m_e}\right)^2 \hat{g}_{ae}$	Primakoff effect, with $\omega \approx m_e$ 0 $-2\omega^2(1 - \cos\theta) = t$ $\frac{2\alpha}{\pi}g_{\alpha e} + o\left(\frac{m_e^2}{\omega^2}\right)$
m _e « m _e	ALP to photon decay	Primakoff effect, with $\omega \gg m_c$	$m_a \ll m_e$	ALP to photon decay	Primakoff effect, with $\omega \gg m_c$	$m_a \ll m_e$	ALP to photon decay	Primakoff effect, with $\omega \gg m_c$
91 q3	0	$-2\omega^2(1 - \cos\theta) = t$	91 92	0	$-2\omega^2(1-\cos\theta) = t$	q_1^2 q_2^2	0	$-2\omega^2(1-\cos\theta) = t$
geff gar	$-\frac{\alpha}{6\pi} \left(\!\frac{m_e}{m_e}\!\right)^2 \hat{g}_{ae}$	$\frac{2\alpha}{\pi}\hat{g}_{\alpha\varepsilon}+O\left(\frac{m_e^2}{\omega^2}\right)$	g_{ay}^{eff}	$-rac{lpha}{6\pi} \left(rac{m_a}{m_e} ight)^2 \hat{g}_{ae}$	$\frac{2\alpha}{\pi}\hat{g}_{ae} + \mathcal{O}\left(\frac{m_e^2}{\omega^2}\right)$	g_{ay}^{eff}	$-rac{lpha}{6\pi} \Big(rac{m_a}{m_e}\Big)^2 \hat{g}_{ae}$	$\frac{2\alpha}{\pi}\hat{g}_{ae} + \mathcal{O}\left(\frac{m_e^2}{\omega^2}\right)$
$\mathbf{m}_e \ll \mathbf{m}_e$	ALP to photon decay	Primakoff effect, with $\omega \gg m_e$	$m_a \ll m_e$	ALP to photon decay	Primakoff effect, with $\omega \gg m_e$	$m_a \ll m_e$	ALP to photon decay	Primakoff effect, with $\omega \gg m_e$
q_{1}^{2}	0	0	q_{1}^{2}	0	0	q_{1}^{2}	0	0
92	0	$-2\omega^2(1-\cos\theta)=t$	q_{2}^{2}	0	$-2\omega^2(1-\cos\theta)=t$	q_{2}^{2}	0	$-2\omega^2(1-\cos\theta)=t$
geff gar	$-\frac{\alpha}{6\pi}\left(\frac{m_a}{m_e}\right)^2 \hat{g}_{ae}$	$\frac{2\alpha}{\pi}\hat{g}_{\alpha\varepsilon} + O\left(\frac{m_{\varepsilon}^2}{\omega^2}\right)$	$g_{a\gamma}^{\mathrm{eff}}$	$-rac{lpha}{6\pi} \left(\!rac{m_a}{m_e}\! ight)^2 \hat{g}_{ae}$	$\frac{2\alpha}{\pi}\hat{g}_{ae} + \mathcal{O}\left(\frac{m_e^2}{\omega^2}\right)$	$g_{a\gamma}^{\mathrm{eff}}$	$-rac{lpha}{6\pi} \left(rac{m_a}{m_e} ight)^2 \widehat{g}_{ae}$	$\frac{2\alpha}{\pi}\hat{g}_{ae} + \mathcal{O}\left(\frac{m_e^2}{\omega^2}\right)$
m _e « m _e	ALP to photon decay	Primakoff effect, with $\omega \gg m_e$	$m_a \ll m_e$	ALP to photon decay	Primakoff effect, with $\omega \gg m_e$	$m_a \ll m_e$	ALP to photon decay	Primakoff effect, with $\omega \gg m_e$
q_{1}^{2}	0	0	q_{1}^{2}	0	0	q_1^2	0	0
			**					
q_{2}^{2}	0	$-2\omega^2(1-\cos\theta)=t$	q_2^2	0	$-2\omega^2(1-\cos\theta)=t$	q_{2}^{2}	0	$-2\omega^2(1-\cos\theta)=t$

Running couplings in aQED

From now on: consider the EFT with $g_{a\gamma}^{\Lambda} \ll \hat{g}_{ae}^{\Lambda}$ i.e. $g_{a\gamma}(\mu) \ll \hat{g}_{ae}(\mu)$

