

Feynman integrals: Turning mathematics into precision predictions

Stefan Weinzierl

May 10, 2023

Section 1

Introduction


Scattering amplitudes

- We would like to make **precise** predictions for observables in scattering experiments from (quantum) field theory.
- Any such calculation will involve a **scattering amplitude**.
- Unfortunately we cannot calculate scattering amplitudes exactly.
- If we have a small parameter like a small coupling, we may use **perturbation theory**.
- We may organise the perturbative expansion of a scattering amplitude in terms of Feynman diagrams.


Scattering amplitude = sum of all Feynman diagrams

Applications


High-energy experiments: LHC




Low-energy experiments: Moller and P2



Gravitational waves:



Spectroscopy: Lamb shift



Standard techniques

- **Dimensional regularisation** ('t Hooft, Veltman '72, Bollini, Giambiagi '72, Ashmore '72):
 $D = 4 - 2\varepsilon$, used to regulate ultraviolet and infrared divergences.
- **Integration-by-parts identities** (Tkachov '81, Chetyrkin '81):
leads to master integrals $I = (I_1, I_2, \dots, I_{N_F})$.
- **Method of differential equations** (Kotikov '90, Remiddi '97, Gehrmann and Remiddi '99):

$$dI = A(x, \varepsilon) I$$

- **Transformation to ε -factorised form** (Henn '13):

$$dI = \varepsilon A(x) I$$

The method of differential equations

We want to calculate

$$I(\varepsilon, x)$$

as a Laurent series in ε .

- 1 Find a differential equation with respect to the kinematic variables for the Feynman integral (*always possible*).
- 2 Transform the differential equation into an ε -factorised form (**bottle neck**).
- 3 Solve the latter differential equation with appropriate boundary conditions (*always possible*).

Example for an ε -factorised form

$$A(x) = C_1 \omega_1 + C_2 \omega_2$$

with differential one-forms

$$\omega_1 = \frac{dx}{x}, \quad \omega_2 = \frac{dx}{x-1},$$

and matrices

$$C_1 = \begin{pmatrix} 2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 \\ 1 & -1 & 0 & 0 & 0 & 2 \end{pmatrix}, \quad C_2 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & -1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -2 \end{pmatrix}.$$

Notation

- $N_F = N_{\text{Fibre}}$: Number of master integrals,
master integrals denoted by $l = (l_1, \dots, l_{N_F})$.
- $N_B = N_{\text{Base}}$: Number of kinematic variables,
kinematic variables denoted by $x = (x_1, \dots, x_{N_B})$.
- $N_L = N_{\text{Letters}}$: Number of letters,
differential one-forms denoted by $\omega = (\omega_1, \dots, \omega_{N_L})$.

- **Fibre** spanned by the master integrals $I = (I_1, \dots, I_{N_F})$.
(The master integrals $I_1(x), \dots, I_{N_F}(x)$ can be viewed as local sections, and for each x they define a basis of the vector space in the fibre.)
- **Base space** with coordinates $x = (x_1, \dots, x_{N_B})$ corresponding to kinematic variables.
- **Connection** defined by the matrix A with differential one-forms $\omega = (\omega_1, \dots, \omega_{N_L})$.

We would like to **transform** this vector bundle **to an ε -factorised form** through

- a change of basis in the fibre,
- a coordinate transformation on the base manifold.

Iterated integrals

Definition

For $\omega_1, \dots, \omega_k$ differential 1-forms on a manifold M and $\gamma: [0, 1] \rightarrow M$ a path, write for the **pull-back** of ω_j to the interval $[0, 1]$

$$f_j(\lambda) d\lambda = \gamma^* \omega_j.$$

The **iterated integral** is defined by

$$I_\gamma(\omega_1, \dots, \omega_k; \lambda) = \int_0^\lambda d\lambda_1 f_1(\lambda_1) \int_0^{\lambda_1} d\lambda_2 f_2(\lambda_2) \dots \int_0^{\lambda_{k-1}} d\lambda_k f_k(\lambda_k).$$

Chen '77

Multiple polylogarithms

Consider differential one-forms on $\mathbb{C} \cup \{\infty\}$ (the **Riemann sphere**) of the form

$$\omega^{\text{mpl}}(z_j) = \frac{d\lambda}{\lambda - z_j}.$$

Definition (Multiple polylogarithms)

$$G(z_1, \dots, z_k; \lambda) = \int_0^\lambda \frac{d\lambda_1}{\lambda_1 - z_1} \int_0^{\lambda_1} \frac{d\lambda_2}{\lambda_2 - z_2} \dots \int_0^{\lambda_{k-1}} \frac{d\lambda_k}{\lambda_k - z_k}, \quad z_k \neq 0$$

Caveats of iterated integrals

- In general, an individual iterated integral is **not homotopy invariant**. The linear combination making up a Feynman integral is, since the connection A is flat (integrable).
- If the differential one-forms ω_k transform nicely under a **group of coordinate transformations**, this does in general not imply that iterated integrals transform nicely as well. However, the vector space spanned by the master integrals does again. Suggests to use different bases of master integrals in different kinematic regions.

Section 2

Geometry

The base space

Question:

After a suitable coordinate transformation, can we relate the base space to a space known from mathematics?

The base space

- Assume we have $(n - 3)$ variables z_1, \dots, z_{n-3} and differential one-forms


$$\omega_k \in \{d\ln(z_1), d\ln(z_2), \dots, d\ln(z_1 - 1), \dots, d\ln(z_i - z_j), \dots\}$$

- The iterated integrals $I_\gamma(\omega_1, \dots, \omega_r; \lambda)$ are **multiple polylogarithms**.
- We require $z_i \notin \{0, 1, \infty\}$ and $z_i \neq z_j$:
This defines the **moduli space** $\mathcal{M}_{0,n}$: The space of configurations of n points on a Riemann sphere modulo Möbius transformations.
- Usually the z_i are functions of the kinematic variables x and the arguments of the dlog-forms define the **Landau singularities**.

Multiple polylogarithms again

Take home message:

Feynman integrals, which evaluate to multiple polylogarithms are related to a Riemann sphere (a smooth complex algebraic curve of genus zero).



Section 3

Elliptic curves

Beyond multiple polylogarithms

- Not every Feynman integral can be expressed in terms of multiple polylogarithms.
- Starting from two-loops, we encounter more complicated functions.
- The next-to-simplest Feynman integrals involve an **elliptic curve**.


Elliptic curves

We do not have to go very far to encounter elliptic integrals in precision calculations: The simplest example is the two-loop electron self-energy in QED:


There are **three Feynman diagrams** contributing to the two-loop electron self-energy in QED with a single fermion:



All master integrals are (sub-) topologies of the **kite graph**:



One sub-topology is the **sunrise graph** with three equal non-zero masses:




(Sabry, '62)

Where is the elliptic curve?

For the sunrise it's very simple: The second graph polynomial defines an elliptic curve in Feynman parameter space:


$$-p^2 a_1 a_2 a_3 + (a_1 + a_2 + a_3)(a_1 a_2 + a_2 a_3 + a_3 a_1) m^2 = 0.$$

Three shades of an elliptic curve




Complex algebraic curve

$$y^2 = 4x^3 - g_2x - g_3$$



Real Riemann surface of
genus one with one
marked point



Complex plane modulo
lattice: \mathbb{C}/Λ


Moduli spaces

$\mathcal{M}_{g,n}$: Space of **isomorphism classes of** smooth (complex, algebraic) **curves of genus g with n marked points.**


complex curve




\Leftrightarrow



real surface



\Leftrightarrow



Coordinates

Genus 0: $\dim \mathcal{M}_{0,n} = n - 3$.

Sphere has a **unique shape**

Use **Möbius transformation** to fix $z_{n-2} = 1, z_{n-1} = \infty, z_n = 0$

Coordinates are **(z_1, \dots, z_{n-3})**

Genus 1: $\dim \mathcal{M}_{1,n} = n$.


One coordinate describes the **shape of the torus**

Use **translation** to fix $z_n = 0$

Coordinates are **$(\tau, z_1, \dots, z_{n-1})$**

Iterated integrals on $\mathcal{M}_{0,n}$ and $\mathcal{M}_{1,n}$

- Iterated integrals on $\mathcal{M}_{0,n}$ with at most simple poles are **multiple polylogarithms**.
Most of the known Feynman integrals fall into this category.
- Iterated integrals on $\mathcal{M}_{1,n}$ are **iterated integrals of modular forms** and **elliptic multiple polylogarithms** (and mixtures thereof).
The simplest example is the two-loop sunrise integral with non-zero masses.



Adams, S.W. '17, Broedel, Duhr, Dulat, Tancredi, '17,
Ch. Bogner, S. Müller-Stach, S.W., '19

Physics is about numbers:

- Iterated integrals of modular forms and elliptic multiple polylogarithms can be evaluated numerically with **arbitrary precision**.
- Implemented in GiNaC.

Walden, S.W, '20

```
ginsh - GiNaC Interactive Shell (GiNaC V1.8.1)
  __, _____ Copyright (C) 1999-2021 Johannes Gutenberg University Mainz,
  (__) *          | Germany. This is free software with ABSOLUTELY NO WARRANTY.
  ._) i N a C | You are welcome to redistribute it under certain conditions.
<-----' For details type `warranty;'.
```

Type ?? for a list of help topics.


```
> Digits=50;
```

```
50
```

```
> iterated_integral({Eisenstein_kernel(3,6,-3,1,1,2)},0.1);
0.23675657575197179243274817775862177623438999192840338805367
```

Generalisations

- We understand by now very well Feynman integrals related to algebraic curves of genus 0 and 1. These correspond to iterated integrals on the moduli spaces $\mathcal{M}_{0,n}$ and $\mathcal{M}_{1,n}$.
- The obvious generalisation is the generalisation to algebraic curves of **higher genus g** , i.e. iterated integrals on the moduli spaces $\mathcal{M}_{g,n}$.
- However, we also need the generalisation from curves to surfaces and **higher dimensional objects**: The geometry of the banana graphs with equal non-vanishing internal masses



are **Calabi-Yau manifolds**.

Section 4

Calabi-Yau manifolds

Definition

A Calabi-Yau manifold of complex dimension n is a compact Kähler manifold M with vanishing first Chern class.

Theorem (conjectured by Calabi, proven by Yau)

An equivalent condition is that M has a Kähler metric with vanishing Ricci curvature.

Mirror symmetry

The **mirror map** relates a Calabi-Yau manifold A to another Calabi-Yau manifold B with Hodge numbers $h_B^{p,q} = h_A^{n-p,q}$.

Candelas, De La Ossa, Green, Parkes '91

			1			
		0		0		
	0		$h^{1,1}$		0	
1		$h^{2,1}$		$h^{2,1}$		1
	0		$h^{1,1}$		0	
		0		0		
			1			

Calabi-Yau manifold A

			1			
		0		0		
	0		$h^{2,1}$		0	
1		$h^{1,1}$		$h^{1,1}$		1
	0		$h^{2,1}$		0	
		0		0		
			1			

mirror image B

Fantastic Beasts and Where to Find Them


- Bananas




- Fishnets




- Amoebas



- Tardigrades



- Paramecia





Aluffi, Marcolli, '09, Bloch, Kerr, Vanhove, '14
Bourjaily, McLeod, von Hippel, Wilhelm, '18
Duhr, Klemm, Loebbert, Nega, Porkert, '22

- The l -loop banana integral with (equal) non-zero masses is related to a **Calabi-Yau $(l-1)$ -fold**.
- An elliptic curve is a Calabi-Yau 1-fold, this is the geometry at two-loops.
- The system of differential equations for the equal mass l -loop banana integral can be transformed to an **ε -factorised form**.
 - Change of variables from $x = p^2/m^2$ to τ given by **mirror map**.
 - Transformation constructed from **special local normal form** of a Calabi-Yau operator.
- Strong support for the conjecture that a transformation to an ε -factorised differential equation exists for all Feynman integrals.

M. Bogner '13, D. van Straten '17

Results: Six loops



Expansion around $y = 0$ converges at six loops for $|p^2| > 49m^2$.

Agrees with results from `pySecDec`.

The geometry of this Feynman integral is a **Calabi-Yau five-fold**.


Pögel, Wang, S.W. '22

Section 5


Phenomenology

Examples

- Dijet production at $N^3\text{LO}$
(related to a Calabi-Yau 2-fold).



- Top pair production at $N^4\text{LO}$
(related to a Calabi-Yau 3-fold)



Conclusions

- Feynman integrals are needed for precision calculations in perturbative quantum field theory.
- Method of differential equations is a powerful tool for computing Feynman integrals.
- Deep connection with mathematics:
 - Vector bundle equipped with a flat connection.
 - Base space related to the moduli space of geometric objects:
Spheres, elliptic curves, Calabi-Yau n -folds, ...