



CP CONSERVATION IN THE STRONG INTERACTIONS

BJÖRN GARBRECHT

Technische Universität München

MITP Mainz

AUGUST 8TH, 2024

The background of the slide is a close-up of red, vertically pleated curtains. The lighting is dramatic, with the center of the curtains being brighter and the edges fading into black.

Collaborators

WEN-YUAN AI

JUAN CRUZ

CARLOS TAMARIT

Outline

- Introduction: QCD θ -parameter, EFT, neutron EDM, topology
- Functional quantization and θ
- Canonical quantization and θ

INTRODUCTION

CP-odd terms in effective field theories

Topology

CP violation in the strong interactions?

No empirical evidence—neutron electric dipole moment (EDM) strongly constrained:

$$d_n = (0.0 \pm 1.1_{\text{stat}} \pm 0.2_{\text{sys}}) \times 10^{-26} e \text{ cm} \quad [2020 \text{ @ PSI}]$$

QCD with massive quarks

$$\mathcal{L} \supset \frac{1}{2g^2} \text{tr} F_{\mu\nu} F^{\mu\nu} + \sum_{j=1}^{N_f} \bar{\psi}_j \left(i\not{\partial} - m_j e^{i\alpha_j \gamma^5} \right) \psi_j + \frac{1}{16\pi^2} \theta \text{tr} F_{\mu\nu} \tilde{F}^{\mu\nu}$$

Believed to cause a neutron electric dipole moment (EDM) $d_n \sim 10^{-15} e \text{ cm} \left(\theta + \sum_j \alpha_j \right)$
[Baluni (1979); Crewther, Di Vecchia, Veneziano, Witten (1979)]

But does it?

CP violation in the strong interactions?

No empirical evidence—neutron electric dipole moment (EDM) strongly constrained:

$$d_n = (0.0 \pm 1.1_{\text{stat}} \pm 0.2_{\text{sys}}) \times 10^{-26} e \text{ cm} \quad [2020 \text{ @ PS}]$$

$\propto \vec{E} \cdot \vec{B}$
parity-odd

$\propto \vec{E}^2 + \vec{B}^2$
 α_j , parity-odd

QCD with massive quarks

$$\mathcal{L} \supset \frac{1}{2g^2} \text{tr} F_{\mu\nu} F^{\mu\nu} + \sum_{j=1}^{N_f} \bar{\psi}_j \left(i\not{\partial} - m_j e^{i\alpha_j \gamma^5} \right) \psi_j + \frac{1}{16\pi^2} \theta \text{tr} F_{\mu\nu} \tilde{F}^{\mu\nu}$$

$\propto \vec{E} \cdot \vec{B}$
parity-odd

Believed to cause a neutron electric dipole moment (EDM) $d_n \sim 10^{-15} e \text{ cm} \left(\theta + \sum_j \alpha_j \right)$
[Baluni (1979); Crewther, Di Vecchia, Veneziano, Witten (1979)]

But does it?

CP violation in the strong interactions?

No empirical evidence—neutron electric dipole moment (EDM) strongly constrained:

$$d_n = (0.0 \pm 1.1_{\text{stat}} \pm 0.2_{\text{sys}}) \times 10^{-26} e \text{ cm} \quad [2020 \text{ @ PSI}]$$

QCD with massive quarks

$$\mathcal{L} \supset \frac{1}{2g^2} \text{tr} F_{\mu\nu} F^{\mu\nu} + \sum_{j=1}^{N_f} \bar{\psi}_j \left(i\not{\partial} - m_j e^{i\alpha_j \gamma^5} \right) \psi_j + \frac{1}{16\pi^2} \theta \text{tr} F_{\mu\nu} \tilde{F}^{\mu\nu}$$

Believed to cause a neutron electric dipole moment (EDM) $d_n \sim 10^{-15} e \text{ cm} \left(\theta + \sum_j \alpha_j \right)$
[Baluni (1979); Crewther, Di Vecchia, Veneziano, Witten (1979)]

But does it?

Effective interactions with θ

$SU(N_f)_L \times SU(N_f)_R$ global symmetry in the limit of massless quarks

Chiral $U(1)_A$ symmetry of the quarks is **anomalous** however

$\longrightarrow \mathcal{L}$ invariant under [Fujikawa (1979,80)]

chiral trafo

$$\begin{aligned}\psi &\rightarrow e^{i\beta\gamma_5}\psi \\ \bar{\psi} &\rightarrow \bar{\psi}e^{i\beta\gamma_5}\end{aligned}$$

plus

“spurion” trafo

$$\begin{aligned}m_j e^{i\alpha_j\gamma_5} &\rightarrow m_j e^{i(\alpha_j - 2N_f\beta)\gamma_5} \\ \theta &\rightarrow \theta + 2N_f\beta\end{aligned}$$

In fact, the “spurions” are those who break the symmetry explicitly.

This pattern should be replicated by any effective theory.

Rephasing invariant: $\bar{\theta} = \theta + \bar{\alpha}$, where $\bar{\alpha} = \sum_{j=1}^{N_f} \alpha_j$, $\longrightarrow \theta$ is an angle

Integrating out gauge fields: Effective 't Hooft vertex

Topological effects described by effective 't Hooft vertex (Γ_{N_f} some coefficient): ['t Hooft (1976,86)]

$$\mathcal{L} + \frac{1}{16\pi^2} \theta \operatorname{tr} F_{\mu\nu} \tilde{F}^{\mu\nu} \rightarrow \mathcal{L} - \Gamma_{N_f} e^{i\xi} \prod_{j=1}^{N_f} (\bar{\psi}_j P_L \psi_j) - \Gamma_{N_f} e^{-i\xi} \prod_{j=1}^{N_f} (\bar{\psi}_j P_R \psi_j)$$

- Effective interaction breaks $U(1)_A$ explicitly $\longrightarrow \eta'$ -mass
- ξ should be expressed in terms of parameters of the fundamental theory
- As a spurion, $\xi \rightarrow \xi + 2N_f\beta$

Two options: $\xi = \theta$ (in general misaligned with masses) $\longrightarrow CP$ violation
 $\xi = -\bar{\alpha}$ (present claim, aligned with mass terms) \longrightarrow no CP violation

So which one is it?

In principle, we could have $\xi = c_\alpha \bar{\alpha} + c_\theta \theta$ for integer $c_{\alpha,\theta}$ (α, θ are angular variables) with $c_\alpha + c_\theta = 1$. We shall see that this general case is not realized in the explicit calculation.

Effective chiral Lagrangian (χ PT)

$$U = U_0 e^{\frac{i}{f_\pi} \Phi} \quad U_0: \text{chiral condensate}$$

$$\Phi = \begin{bmatrix} \pi^0 + \frac{1}{\sqrt{3}}\eta + \sqrt{\frac{2}{3}}\eta' & \sqrt{2}\pi^+ & \sqrt{2}K^+ \\ \sqrt{2}\pi^- & -\pi^0 + \frac{1}{\sqrt{3}}\eta + \sqrt{\frac{2}{3}}\eta' & \sqrt{2}K^0 \\ \sqrt{2}K^- & \sqrt{2}\bar{K}^0 & -\frac{2}{\sqrt{3}}\eta + \sqrt{\frac{2}{3}}\eta' \end{bmatrix}$$

Chiral Lagrangian (lowest order terms) inherits “spurious” symmetries:

$$\mathcal{L} = \frac{f_\pi^2}{4} \text{Tr} \partial_\mu U \partial^\mu U^\dagger + \frac{f_\pi^2 B_0}{2} \text{Tr}(MU + U^\dagger M^\dagger) + |\lambda| e^{-i\xi} f_\pi^4 \det U + |\lambda| e^{i\xi} f_\pi^4 \det U^\dagger \\ + i\bar{N}\not{\partial}N - \left(m_N \bar{N} \tilde{U} P_L N + ic \bar{N} \tilde{U}^\dagger \not{\partial} P_L \tilde{U} N + d \bar{N} \tilde{M}^\dagger P_L N + e \bar{N} \tilde{U} \tilde{M} \tilde{U} P_L N + \text{h.c.} \right)$$

$$M = \text{diag}\{m_u e^{i\alpha_u}, m_d e^{i\alpha_d}, m_s e^{i\alpha_s}\} \quad \text{nucleon doublet } N = \begin{pmatrix} p \\ n \end{pmatrix} \\ \tilde{M}, \tilde{U} \text{ reduced to subspace } (u, d)$$

Effective interaction $\propto \det U$ cannot be quantitatively reliably handled in χ PT but yet represents pattern of broken axial symmetry.

CP -odd neutron interactions [e.g. Srednicki QFT (2007)]

- Write $U_0 = \langle U \rangle = \text{diag}(e^{i\varphi_u}, e^{i\varphi_d}, e^{i\varphi_s})$
- Minimize $V(\langle U \rangle) \rightarrow m_i(\varphi_i + \alpha_i) \approx \bar{m}(m_u, m_d, m_s)(\xi + \alpha_u + \alpha_d + \alpha_s)$
(for small angles)
- Substitute φ_i back into \mathcal{L} & suitably redefine $N \rightarrow \mathcal{N}(N, U)$

$$\mathcal{L}_{\text{neutron}} \supset -\frac{2c+1}{f_\pi} \partial_\mu \pi^a \bar{N} T^a \gamma^\mu \gamma_5 \mathcal{N} \quad CP \text{ even}$$
$$+ \frac{2(d+e)\bar{m}}{f_\pi} (\xi + \alpha_u + \alpha_d + \alpha_s) \bar{N} \pi^a T^a \mathcal{N} \quad CP \text{ odd}$$

Neutron electric dipole moment

$$i\mathcal{M} = -2iD(q^2)\epsilon_{\mu}^*(\vec{q})\bar{u}_{s'}(\vec{p}')\frac{i}{4}[\gamma^{\mu},\gamma^{\nu}]q_{\nu}i\gamma_5u_s(\vec{p})$$

$$\rightarrow \mathcal{L}_{\text{eff}} = D(0)\bar{n}\underbrace{F_{\mu\nu}\frac{i}{4}[\gamma^{\mu},\gamma^{\nu}]i\gamma_5}_{\subset \vec{S}\cdot\vec{E}}n$$



- χ PT value: $d_n = 3.2 \times 10^{-16}(\xi + \bar{\alpha})e \text{ cm}$
- Experimental bound: $|d_n| < 1.8 \times 10^{-26}e \text{ cm}$ (90% c.l.) [nEDM/PSI (2020)]
- Calculations e.g. of neutron EDM implicitly *assume* $\xi = \theta$
[e.g. Baluni (1979); Crewther, Di Vecchia, Veneziano, Witten (1979)]
- However $\xi = -\bar{\alpha}$ also perfectly valid by arguments used to this end
- Another signature—weaker bounds: $\eta' \rightarrow \pi\pi$

Topology in four-dimensional spacetime—winding number Δn

$$U = \begin{pmatrix} a_R + ia_I & -b_R + ib_I \\ b_R + ib_I & a_R - ia_I \end{pmatrix} \in \text{SU}(2) \text{ for } a_R^2 + a_I^2 + b_R^2 + b_I^2 = 1$$

$$\Rightarrow \text{Homotopy: } \text{SU}(3) \supset \text{SU}(2) \cong S^3 \longrightarrow \pi_3(\text{SU}(2)) = \pi_3(S^3) = \mathbb{Z}$$

Theta-term/topological term is a total divergence:

gauge
invariant

$$\frac{1}{4} \text{tr} F_{\mu\nu} \tilde{F}_{\mu\nu} = \partial_\mu K_\mu$$

$$K_\mu = \epsilon_{\mu\nu\alpha\beta} \text{tr} \left[\frac{1}{2} A_\nu \partial_\alpha A_\beta + \frac{1}{3} A_\nu A_\alpha A_\beta \right]$$

gauge
dependent


Topological quantization for pure gauge $A_\mu \rightarrow -\frac{i}{g}(\partial_\mu U)U^{-1}$ at $\partial\Omega \cong S^3$

$$\Delta n = \frac{1}{16\pi^2} \int_\Omega d^4x F_{\mu\nu} \tilde{F}_{\mu\nu} = \frac{1}{4\pi^2} \oint_{\partial\Omega} d^3\sigma K_\perp \in \mathbb{Z}$$

Haar measure for pure gauge

$$K_\mu = \frac{1}{6} \epsilon_{\mu\nu\lambda\rho} \text{tr}[(U^{-1}\partial_\nu U)(U^{-1}\partial_\lambda U)(U^{-1}\partial_\rho U)]$$

E.g. take boundary of $\Omega = \mathbb{R}^4$ as a sphere S^3 :



Or $\Omega = T^4$ (lattice), $\Omega = S^4$ (Euclidean dS): $\Delta n \in \mathbb{Z}$ based on slightly more involved argument

Topology—instantons


Does $\Delta n \neq 0$ imply nontrivial physical field configurations?

Yes, cf. anti-instanton: $A_\mu^u{}_\nu = -\frac{\sigma_{\mu\nu}^u x_\nu}{x^2 + \rho^2}$

(extended solution to *Euclidean* EOMs)

[Belavin, Polyakov, Schwarz, Tyupkin (1975)]

Surface term decays as $1/|x|^3 \rightarrow$ surface integral does not need to vanish



Theta term contributes to the action though being a total derivative

Topology on spatial hypersurfaces—point compactification, large gauge transformations

Consider temporal gauge $A^0 = 0$ (in view of canonical quantization)

Chern–Simons functional:


$$W[\vec{A}] = \frac{1}{4\pi^2} \epsilon_{ijk} \int_V d^3x \operatorname{tr} \left[\frac{1}{2} A_i \partial_j A_k - \frac{i}{3} A_i A_j A_k \right] \equiv \frac{1}{4\pi^2} \int_V d^3x K_0$$

Define $\vec{A}_U = U \vec{A} U^{-1} + i U^{-1} \vec{\nabla} U$ (residual gauge freedom in temporal gauge)

Assume $\vec{A} = i U^{-1} \vec{\nabla} U$ (i.e. pure gauge) on ∂V

With extra constraint $U(\vec{x}) \rightarrow \text{const.}$ on ∂V (periodic on T^3)

→ Point compactification, homotopy $V \cong S^3$ ($V \cong T^3$)



$U^{(n)}$: “large” ($n \neq 0$) gauge transformation on spacelike ($\tau = \text{const.}$) hypersurface $V \simeq S^3$ ($V \simeq T^3$) changing the Chern–Simons number by $n = W[\vec{A}_{U^{(n)}}] - W[\vec{A}] \in \mathbb{Z}$ units

Topology on spatial hypersurfaces—point compactification, large gauge transformations

Consider temporal gauge $A^0 = 0$ (in view of canonical quantization)

Chern–Simons

Equivalence classes of $U^{(n)}$ (not connected with $\mathbb{1}$ for $n \neq 0$) only exist when we require these added *constraints* on $\vec{A}(\vec{x})$ (beyond $A^0(\vec{x}) = 0$)

W

$x K_0$

Cannot impose these unless properly taken account in canonical formalism

Define $\vec{A}_U =$

(gauge)

Assume $\vec{A} = i$

Unlike Δn , the equivalence classes are a result of a gauge choice.

With extra constraint $U(x) \rightarrow \text{const.}$ on ∂V (periodic on T^3)

space

$A = iU^{-1} \vec{\nabla} U$
& $U(\vec{x}) = \text{const.}$
on boundary



(three)-sphere

→ Point compactification, homotopy $V \cong S^3$ ($V \cong T^3$)

$U^{(n)}$: “large” ($n \neq 0$) gauge transformation on spacelike ($\tau = \text{const.}$) hypersurface $V \simeq S^3$ ($V \simeq T^3$) changing the Chern–Simons number by $n = W[\vec{A}_{U^{(n)}}] - W[\vec{A}] \in \mathbb{Z}$ units

FUNCTIONAL QUANTIZATION

Theta in infinite spacetime volume

Why $T \rightarrow \infty$

(Implying $\Omega = VT \rightarrow \infty$ as opposed to a finite spacetime volume)

To evaluate amplitudes at finite T , project path integral on the state in terms of a wave function(al) $\Psi[\phi(\vec{x})] = \langle \phi(\vec{x}) | \Psi \rangle$ (more on this later):

$$\langle \Psi_f, t_f | \Psi_i, t_i \rangle = \int \mathcal{D}\phi_f \mathcal{D}\phi_i \langle \Psi_f, t_f | \phi_f \rangle \int \mathcal{D}\phi e^{iS[\phi]} \langle \phi_i | \Psi_i, t_i \rangle$$

$|\phi_{i,f}\rangle$ field eigenstates, not energy eigenstates

$$\begin{aligned} \phi(t_f, \vec{x}) &= \phi_f(\vec{x}) \\ \phi(t_i, \vec{x}) &= \phi_i(\vec{x}) \end{aligned}$$

Problem: Neither know $\Psi[\phi(\vec{x})]$ nor kernels of Schrödinger equation

Way out: Euclidean path integral/project on ground state

$$\lim_{T \rightarrow \infty} \frac{e^{-HT}}{e^{-E_0 T}}$$

or

$$\lim_{T \rightarrow \infty} \frac{e^{-iHT(1-i\epsilon)}}{e^{-iE_0 T(1-i\epsilon)}}$$

H : Hamiltonian

E_0 : ground state energy

→ Obtain vacuum correlations without bothering about $\Psi[\phi(\vec{x})]$

Boundary configurations & topological quantization

The parameter θ can be viewed as an angular variable
(forced by the anomalous chiral current). \rightarrow

Requires $\Delta n \in \mathbb{Z}$ (“topological quantization”) $\rightarrow \exp(iS)|_{\theta} = \exp(iS)|_{\theta+2\pi}$

Readily built into the path integral for $VT \rightarrow \infty$ without constraining boundary conditions by hand:

(Relatively) nonvanishing contributions in *infinite* spacetime only from classical local minima of the Euclidean action & fluctuations about these—these configurations must go to pure gauges at ∞

There is no such restriction/principle to fixed physical bcs. for finite VT .


Indeed, for pure gauge configurations at $\infty \rightarrow \Delta n \in \mathbb{Z}$ (as discussed above)

Consequence: $\Delta n \in \mathbb{Z}$ requires $T \rightarrow \infty$ first \rightarrow in the path integral, take $T \rightarrow \infty$, then sum over all topological sectors Δn weighted $\exp(i\Delta n\theta)$

More technically: Integration contour from Lefschetz thimbles

Parametrization of the path integral through steepest descent contours about classical saddle points \rightarrow Contour integration on Lefschetz thimbles

$$\frac{\partial \phi(x; u)}{\partial u} = \frac{\overline{\delta S_{\mathbb{E}}[\phi(x; u)]}}{\delta \phi(x; u)} \implies -\frac{\partial \text{Re} S_{\mathbb{E}}[\phi(x; u)]}{\partial u} \leq 0 \quad \text{and} \quad \frac{\partial \text{Im} S_{\mathbb{E}}[\phi(x; u)]}{\partial u} = 0$$



Each thimble emerges from a critical point and corresponds to one $\Delta n \in \mathbb{Z}$

Keeping VT finite while summing over different Δn does *not* correspond to a nonsingular deformation of the contour

Integration contour sweeps over full thimbles first:

$$Z = \lim_{N \rightarrow \infty} \sum_{\Delta n = -N}^N \lim_{VT \rightarrow \infty} \int_{\Delta n} \mathcal{D}\phi e^{-S_{\mathbb{E}}[\phi]}$$

So is it $\xi = -\bar{\alpha}$ or $\xi = \theta$?

The effective vertex generates the following correlation functions at tree level:

$$\left\langle \prod_{j=1}^{N_f} \psi_j(x_j) \bar{\psi}_j(x'_j) \right\rangle_{\text{inst}} = \left(e^{-i\xi} \prod_{j=1}^{N_f} P_{Lj} + e^{i\xi} \prod_{j=1}^{N_f} P_{Rj} \right) \bar{H}(x_1, \dots, x'_1, \dots)$$

Goal: Compute correlation function and compare with EFT answer above to fix ξ

Cf. leading contribution to two-point function

$$\langle \psi_i(x) \psi_j(x') \rangle = i S_{0\text{inst } ij}(x, x')$$

$$i S_{0\text{inst } ij}(x, x') = (-\gamma^\mu \partial_\mu + i m_i e^{-i\alpha_i \gamma^5}) \int \frac{d^4 p}{(2\pi)^4} e^{-ip(x-x')} \frac{\delta_{ij}}{p^2 - m_i^2 + i\epsilon}$$

So $\xi = \theta/\xi = -\bar{\alpha}$ implies CP -violation/no CP -violation

Only one explicit calculation based on dilute instanton gas (DIGA) finding $\xi = \theta$

[t Hooft (1986)]

Fermion correlations

- Obtain correlation functions from Green's functions in fixed background of instantons and anti-instantons
- Interfere all instanton configurations
 - First, within one topological sector
 - Then over the different sectors

Green's function in n -instanton, \bar{n} -anti-instanton background (DIGA)

$$iS_{n,\bar{n}}(x, x') \approx iS_{0\text{inst}}(x, x') + \sum_{\bar{\nu}=1}^{\bar{n}} \frac{\hat{\psi}_{0L}(x - x_{0,\bar{\nu}})\hat{\psi}_{0L}^\dagger(x' - x_{0,\bar{\nu}})}{m e^{-i\alpha}} + \sum_{\nu=1}^n \frac{\hat{\psi}_{0R}(x - x_{0,\nu})\hat{\psi}_{0R}^\dagger(x' - x_{0,\nu})}{m e^{i\alpha}}$$

$\hat{\psi}_{0L,R}$: 't Hooft zero modes

Comments:

- For small masses, zero modes dominate close to the cores of instantons, far away from instantons the solution goes to the zero-instanton configuration [Diakonov, Petrov (1986)]
- **Alignment** of phase α between Lagrangian mass and instanton-induced χSB \rightarrow No indication of CP violation here

Fermion correlations

- Obtain correlation functions from Green's functions in fixed background of instantons and anti-instantons
- Interfere all instanton configurations
 - First, within one topological sector
 - Then over the different sectors

Green's function in n -instanton, \bar{n} -anti-instanton background (DIGA)

$$iS_{n,\bar{n}}(x, x') \approx iS_{0\text{inst}}(x, x') + \sum^{\bar{n}} \frac{\hat{\psi}_{0L}(x - x_{0,\bar{\nu}}) \hat{\psi}_{0L}^\dagger(x' - x_{0,\bar{\nu}})}{m e^{-i\alpha}} + \sum^n \frac{\hat{\psi}_{0R}(x - x_{0,\nu}) \hat{\psi}_{0R}^\dagger(x' - x_{0,\nu})}{m e^{i\alpha}}$$

cf.

$$iS_{0\text{inst}}(x, x') = (-\gamma^\mu \partial_\mu + i m e^{-i\alpha \gamma^5}) \int \frac{d^4 p}{(2\pi)^4} e^{-ip(x-x')} \frac{1}{p^2 - m^2 + i\epsilon}$$

- For small masses, zero modes dominate close to the cores of instantons, far away from instantons the solution goes to the zero-instanton configuration [Diakonov, Petrov (1986)]
- **Alignment** of phase α between Lagrangian mass and instanton-induced χ SB \rightarrow No indication of CP violation here

Fermion correlations

- Obtain correlation functions from Green's functions in fixed background of instantons and anti-instantons
- Interfere all instanton configurations
 - First, within one topological sector
 - Then over the different sectors

Green's function in n -instanton, \bar{n} -anti-instanton background (DIGA)

$$iS_{n,\bar{n}}(x, x') \approx iS_{0\text{inst}}(x, x') + \sum_{\bar{\nu}=1}^{\bar{n}} \frac{\hat{\psi}_{0L}(x - x_{0,\bar{\nu}})\hat{\psi}_{0L}^\dagger(x' - x_{0,\bar{\nu}})}{m e^{-i\alpha}} + \sum_{\nu=1}^n \frac{\hat{\psi}_{0R}(x - x_{0,\nu})\hat{\psi}_{0R}^\dagger(x' - x_{0,\nu})}{m e^{i\alpha}}$$

$\hat{\psi}_{0L,R}$: 't Hooft zero modes

Comments:

- For small masses, zero modes dominate close to the cores of instantons, far away from instantons the solution goes to the zero-instanton configuration [Diakonov, Petrov (1986)]
- **Alignment** of phase α between Lagrangian mass and instanton-induced χSB \rightarrow No indication of CP violation here

Interferences within the topological sectors

Within a topological sector, interfere/sum/integrate over

- all instanton/anti-instanton numbers $n + \bar{n}$ with $\Delta n = n - \bar{n}$ fixed
- locations of all instantons/anti-instantons
- remaining collective coordinates

→ Dilute instanton gas approximation (skip technicalities)

Can also obtain coincident fermion correlations using the index theorem and anomalous current only

Correlation function for fixed Δn

$$\begin{aligned}
 \langle \psi(x) \bar{\psi}(x') \rangle_{\Delta n} &= \sum_{\substack{\bar{n}, n \geq 0 \\ n - \bar{n} = \Delta n}} \frac{1}{\bar{n}! n!} \left[\bar{h}(x, x') \left(\frac{\bar{n}}{m e^{-i\alpha}} P_L + \frac{n}{m e^{i\alpha}} P_R \right) (VT)^{\bar{n}+n-1} + i S_{0\text{inst}}(x, x') (VT)^{\bar{n}+n} \right] \\
 &\quad \times (i\kappa)^{\bar{n}+n} (-1)^{n+\bar{n}} e^{i\Delta n(\alpha+\theta)} \\
 &= \left[(e^{i\alpha} I_{\Delta n+1}(2i\kappa VT) P_L + e^{-i\alpha} I_{\Delta n-1}(2i\kappa VT) P_R) \frac{i\kappa}{m} \bar{h}(x, x') + I_{\Delta n}(2i\kappa VT) i S_{0\text{inst}}(x, x') \right] \\
 &\quad \times (-1)^{\Delta n} e^{i\Delta n(\alpha+\theta)}
 \end{aligned}$$

Instantons per spacetime volume: $i\kappa \propto e^{-S_E}$

χ SB rank-two spinor-tensor from integrating quark zero-modes over the locations of the instantons: $\bar{h}(x, x')$

Modified Bessel function: $I_\nu(x)$

Sum is dominated by particular value of $n \approx \bar{n}$: [Diakonov, Petrov (1986)]

$$\langle n \rangle = \frac{\sum_{n=0}^{\infty} n \frac{(\kappa VT)^n}{n!}}{\sum_{n=0}^{\infty} \frac{(\kappa VT)^n}{n!}} = \kappa VT, \quad \frac{\sqrt{\langle (n - \langle n \rangle)^2 \rangle}}{\langle n \rangle} = \frac{1}{\sqrt{\kappa VT}}, \quad \text{cf. } \lim_{x \rightarrow \infty} \frac{I_{\Delta n}(ix e^{-i0^+})}{I_{\Delta n'}(ix e^{-i0^+})} = 1$$

→ No relative CP phase between mass and instanton induced breaking of χ ral symmetry—**alignment** in infinite-volume limit

Correspondingly, partition function for fixed Δn : [cf. Leutwyler, Smilga (1992)]

$$Z_{\Delta n} = I_{\Delta n}(2i\kappa VT) (-1)^{\Delta n} e^{i\Delta n(\alpha+\theta)}$$

Note: The topological phase $e^{i\Delta n(\alpha+\theta)}$ multiplies $\langle \psi(x)\bar{\psi}(x') \rangle_{\Delta n}$ and $Z_{\Delta n}$ entirely—not just the contributions induced by instantons.

Other correlation functions (n point, stress-energy, for some observer,...) are calculated from the Feynman diagram with the Green's function in the n instanton, \bar{n} anti-instanton background.

Then it remains to average over n , \bar{n} , locations and remaining collective coordinates.

There is no CP violation/misalignment of phases to this end. It remains to consider the interference between the topological sectors.

Interferences among topological sectors (are immaterial)

Topological quantization \leftrightarrow Interference between sectors for $VT \rightarrow \infty$

Fermion correlator

$$\begin{aligned}\langle \psi(x) \bar{\psi}(x') \rangle &= \lim_{\substack{N \rightarrow \infty \\ N \in \mathbb{N}}} \lim_{VT \rightarrow \infty} \frac{\sum_{\Delta n = -N}^N \langle \psi(x) \bar{\psi}(x') \rangle_{\Delta n}}{\sum_{\Delta n = -N}^N Z_{\Delta n}} \\ &= iS_{0\text{inst}}(x, x') + i\kappa \bar{h}(x, x') m^{-1} e^{-i\alpha\gamma^5} \quad (\text{same as for fixed } \Delta n)\end{aligned}$$

Recall: $iS_{0\text{inst}}(x, x') = (-\gamma^\mu \partial_\mu + im e^{-i\alpha\gamma^5}) \int \frac{d^4 p}{(2\pi)^4} e^{-ip(x-x')} \frac{1}{p^2 - m^2 + i\epsilon}$

\longrightarrow No relative CP -phase between mass and instanton term

$$\longrightarrow \xi = -\alpha$$

$\longrightarrow CP$ is conserved

Limits ordered the other way around

First sum over all Δn as well:

$$\begin{aligned} & \sum_{\bar{n}, n \geq 0} \frac{1}{\bar{n}! n!} \left[\bar{h}(x, x') (\bar{n} m^{-1} e^{i\alpha} P_L + n m^{-1} e^{-i\alpha} P_R) (VT)^{\bar{n}+n-1} + iS_{0\text{inst}}(x, x') (VT)^{\bar{n}+n} \right] \\ & \qquad \qquad \qquad \times (-mi\kappa)^{\bar{n}+n} e^{i\Delta n(\alpha+\theta)} \\ & = \left[- (e^{-i\theta} P_L + e^{i\theta} P_R) \frac{i\kappa}{m} \bar{h}(x, x') + iS_{0\text{inst}}(x, x') \right] e^{-2i\kappa VT \cos(\alpha+\theta)} \end{aligned}$$

$$Z \rightarrow \sum_{n, \bar{n}} \frac{1}{n! \bar{n}!} (-i\kappa VT)^{\bar{n}+n} e^{-i(\bar{n}-n)(\alpha+\theta)} = e^{-2i\kappa VT \cos(\alpha+\theta)}$$

Then, $VT \rightarrow \infty$ trivial as VT -dependence cancels

→ Relative CP phase leading to CP -violating observables

However: Changing the order does not correspond to a nonsingular integration contour.

Reduced argument without instantons

- Take $\langle F(x)\tilde{F}(x) \rangle$ as measure for CP violation
- Each element in the sequence over N vanishes (not so when limits ordered the other way around):

$$\langle F(x)\tilde{F}(x) \rangle = \lim_{\substack{N \rightarrow \infty \\ N \in \mathbb{N}}} \lim_{VT \rightarrow \infty} \frac{\sum_{\Delta n=-N}^N \frac{\Delta n}{VT} Z_{\Delta n}}{\sum_{\Delta n=-N}^N Z_{\Delta n}} = 0$$

- Index theorem: No L/R imbalance in fermion zero modes \rightarrow Zero modes remain aligned with quark mass after interference of Δn -sectors

CANONICAL QUANTIZATION
(in finite and infinite volumes)

Theta vacuum, standard story

Take, $A^0 = 0$, *assume* in addition:

For $|\vec{x}| \rightarrow \infty$: $\vec{A}(\vec{x}) = iU^{-1}(\vec{x})\vec{\nabla}U(\vec{x})$ and $U(\vec{x}) \rightarrow \text{const.}$


But why? [cf. Jackiw (1980)]

Consider initial and final states, taking $x_4 \rightarrow \pm\infty$

→ Construct from pure gauge configurations on these surfaces, with

$$\Delta n = \frac{1}{16\pi^2} \int d^4x F_{\mu\nu} \tilde{F}_{\mu\nu} = n_\infty - n_{-\infty} \quad \text{gauge invariant}$$

$$n_{\pm\infty} = \frac{1}{4\pi^2} \int_{x^4=\pm\infty} d^3\sigma K_\perp \in \mathbb{Z} \quad \begin{array}{l} \text{Chern-Simons number} \\ \text{not gauge invariant} \end{array} \quad \begin{array}{l} \text{point com-} \\ \text{pactification} \end{array}$$



Gauge transformations change $n_{\pm\infty}$ by same number of integer units

Construct ground states from prevacua: $n_{-\infty} \rightarrow |n\rangle$ (field eigenstates)
 $n_\infty \rightarrow \langle n|$

Gauge invariant (up to phase) state $|\theta\rangle = \sum_n e^{-in\theta} |n\rangle$

[Callan, Dashen, Gross (1976);
 Jackiw, Rebbi (1976); Jackiw (1980)]

Standard story: two loose ends

The prevacua $|n\rangle$ are field eigenstates, very different from the ground state

Resolutions:

- Take $T \rightarrow \infty$ in the path integral to project on the ground state:
 $|\text{vac}\rangle = e^{-HT} \sum_n e^{-in\theta} |n\rangle$, $T \rightarrow \infty$ (cf. $VT \rightarrow \infty$ in previous part)
- Or use the symmetries and no further properties of the wave functional
[Jackiw, Rebbi (1976); Jackiw (1980)]

States are not normalizable in the proper sense because $\langle \theta^{(i)} | \theta^{(j)} \rangle = \delta(\theta^{(i)} - \theta^{(j)})$

[cf. e.g. Callan, Dashen, Gross (1976); issue taken by Okubo, Marshak (1992)]

Without ado, this contradicts 1st postulate of quantum mechanics.

Possible resolutions:

- Construct wave packets—not acceptable however because gauge invariance should be exact
- Use gauge fixing in order to normalize states (which is what we will do here)

The prevacua $|n\rangle$

Resolutions:

- Take $T \rightarrow \infty$
 $|\text{vac}\rangle = e^{-HT}$

- Or use the sy:
[Jackiw, Rebbi (1976)]

States are not nor:

[cf. e.g. Callan, Dashen, Gr

Without ado, this

Possible resolutions:

- Construct wave packets—not acceptable however because gauge invariance should be exact
- Use gauge fixing in order to normalize states (which is what we will do here)

B. STATEMENT OF THE POSTULATES

1. Description of the state of a system

In chapter I, we introduced the concept of the quantum state of a particle. We first characterized this state at a given time by a square-integrable wave function. Then, in chapter II, we associated a ket of the state space \mathcal{E}_r with each wave function: choosing $|\psi\rangle$ belonging to \mathcal{E}_r is equivalent to choosing the corresponding function $\psi(\mathbf{r}) = \langle \mathbf{r} | \psi \rangle$. Therefore, the quantum state of a particle at a fixed time is characterized by a ket of the space \mathcal{E}_r . In this form, the concept of a state can be generalized to any physical system.

First Postulate: At a fixed time t_0 , the state of a physical system is defined by specifying a ket $|\psi(t_0)\rangle$ belonging to the state space \mathcal{E} .

[Cohen-Tanoudji, Diu, Laloë]

Standard story: two loose ends

The prevacua $|n\rangle$ are field eigenstates, very different from the ground state

Resolutions:

- Take $T \rightarrow \infty$ in the path integral to project on the ground state:
 $|\text{vac}\rangle = e^{-HT} \sum_n e^{-in\theta} |n\rangle$, $T \rightarrow \infty$ (cf. $VT \rightarrow \infty$ in previous part)
- Or use the symmetries and no further properties of the wave functional
[Jackiw, Rebbi (1976); Jackiw (1980)]

States are not normalizable in the proper sense because $\langle \theta^{(i)} | \theta^{(j)} \rangle = \delta(\theta^{(i)} - \theta^{(j)})$

[cf. e.g. Callan, Dashen, Gross (1976); issue taken by Okubo, Marshak (1992)]

Without ado, this contradicts 1st postulate of quantum mechanics.

Possible resolutions:

- Construct wave packets—not acceptable however because gauge invariance should be exact
- Use gauge fixing in order to normalize states (which is what we will do here)

Canonical quantization of the gauge field

Minkowski spacetime, temporal gauge $A^0 = 0$, no sources \rightarrow

$$g\vec{E}^a = -\partial/\partial t \vec{A}^a$$

$$g\vec{B}^a = \vec{\nabla} \times \vec{A}^a - 1/2 f^{abc} \vec{A}^a \times \vec{A}^b$$

Canonical momentum conjugate to \vec{A}^a :

$$g\vec{\Pi}^a = -\vec{E}^a + \frac{g^2}{8\pi^2} \theta \vec{B}^a$$

The corresponding operator must observe the commutation relations:

$$[A^{a,i}(\vec{x}), \Pi^{b,j}(\vec{x}')] = i\delta^{ij} \delta^{ab} \delta^3(\vec{x} - \vec{x}'), \quad [\Pi^{a,i}(\vec{x}), \Pi^{b,j}(\vec{x}')] = 0$$

These commutators hold for (θ_Π arbitrary) $\vec{\Pi}^a = \frac{\delta}{i\delta \vec{A}^a} + \theta_\Pi \frac{g}{8\pi^2} \vec{B}^a$

Hamiltonian density:

$$\mathcal{H} = \frac{1}{2} \left((\vec{E}^a)^2 + (\vec{B}^a)^2 \right) = \frac{1}{2} \left(\left(g \frac{\delta}{i\delta \vec{A}^a} - \frac{g^2}{8\pi^2} (\theta - \theta_\Pi) \vec{B}^a \right)^2 + (\vec{B}^a)^2 \right)$$

No constraints on ∂V accounted for
 $\Psi[\vec{A}]$ must be defined for $U(\vec{x}) \neq \text{const.}$ on ∂V
Residual gauge dofs.: Throw out unphysical states
"First quantize, then constrain"

Wave functional in gauge theory (temporal gauge $A_0 = 0$)

Since $[W[\vec{A}], H] = 0$, can find states $\Psi_{\theta^{(i)}}[\vec{A}]_{(U_4^{(1)})^n} = e^{in\theta^{(i)}} \Psi_{\theta^{(i)}}[\vec{A}]$

Wave functionals not properly normalizable

$$\begin{aligned} \int \mathcal{D}\vec{A} \Psi_{\theta^{(i)}}^{(a)*}[\vec{A}] \Psi_{\theta^{(j)}}^{(b)}[\vec{A}] &= \sum_{\nu=-\infty}^{\infty} \int_{0 \leq W[\vec{A}] < 1} \mathcal{D}\vec{A} e^{-i(\theta^{(i)} - \theta^{(j)})(W[\vec{A}] + \nu)} \psi_{\theta^{(i)}}^{(a)*}[\vec{A}] \psi_{\theta^{(j)}}^{(b)}[\vec{A}] \\ &= 2\pi \delta(\theta^{(i)} - \theta^{(j)}) \int_{0 \leq W[\vec{A}] < 1} \mathcal{D}\vec{A} e^{-i(\theta^{(i)} - \theta^{(j)})W[\vec{A}]} \psi_{\theta^{(i)}}^{(a)*}[\vec{A}] \psi_{\theta^{(j)}}^{(b)}[\vec{A}] \\ &= 2\pi \delta(\theta^{(i)} - \theta^{(j)}) \delta_{ab} \end{aligned}$$

Cf. T^4 /lattice:

$$Z = \prod_a \int \mathcal{D}\vec{A} \Psi_{\theta^{(i)}}^{(a)*}[\vec{A}] e^{-\beta H} \Psi_{\theta^{(i)}}^{(b)}[\vec{A}]$$


Not properly normalizable either

Crystal or circle?

The functionals $\Psi_\theta(\vec{A})$ with above periodicity properties can be viewed as Bloch states.

Bloch states live on a crystal:

$\vec{A}_{U_4^{(1)}}$ is a *different* site than \vec{A}




In contrast: In gauge theory

$\vec{A}_{U_4^{(1)}}$ is a *redundant*

description of the configuration

\vec{A} —corresponding to

$\varphi \rightarrow \varphi + 2\pi n$ on a circle



On a crystal: Bloch states do not correspond to normalized wave functions, these are rather wave packets made up of Bloch states. Packets, however, not translation (gauge) invariant

On a circle: Truncation of the inner product according to a single period leads to properly normalizable states, corresponding here to *gauge fixing* $\vec{A} \in \mathcal{A}$ so that each physical configuration appears one time and one time only:

$$\int_{\mathcal{A}} \underbrace{\mathcal{D}\vec{A} f_{\mathcal{A}}[\vec{A}]}_{\text{gauge invariant under change of } \mathcal{A}} \Psi_{\theta^{(i)}}^{(a)*}[\vec{A}] \Psi_{\theta^{(j)}}^{(b)}[\vec{A}]$$

Note: Under gauge fixed inner product, $\Psi_{\theta^{(i)}}^{(a)}$, $\Psi_{\theta^{(j)}}^{(b)}$ no longer orthogonal for $\theta^{(i)} \neq \theta^{(j)}$

Form of the wave functional

Require: $\frac{\delta}{i\delta\vec{A}(\vec{x})}$ should remain Hermitian under restricted inner product

$\implies \Psi^{(a)}[\vec{A}] \stackrel{(*)}{=} \Psi^{(a)}[\vec{A}]_{\text{g.i.}} \exp(i\varphi[\vec{A}])$ valid for *all* $U(\vec{x})$ (also nonconstant on boundary)

gauge invariant ↗

↖ independent of state (a)

Problem: $\int d^3x \text{tr} \vec{B} \cdot \frac{\delta}{i\delta\vec{A}}$ leads to dependence of pure gauge on other directions

→ “Diagonalize” H : $\Psi'[\vec{A}] = e^{-i(\theta-\theta_\Pi)W[\vec{A}]} \Psi[\vec{A}]$,

$$\frac{\delta}{\delta\vec{A}(\vec{x})} W[\vec{A}] = \frac{g}{8\pi^2} \vec{B}(\vec{x})$$

$$H' = e^{-i(\theta-\theta_\Pi)W[\vec{A}]} H e^{i(\theta-\theta_\Pi)W[\vec{A}]} = \frac{1}{2} \int d^3x \text{tr} \left[-g^2 \frac{\delta^2}{\delta\vec{A}^2} + \vec{B}^2 \right]$$

$$= -\frac{g^2}{2} \int_{\sigma} \frac{\delta^2}{\delta A^2(\sigma)} + \frac{1}{2} \int d^3x \text{tr} \vec{B}^2, \quad \sigma \in \{\sigma_{\text{gauge}}, \sigma_{\parallel}\}$$

Only trivial one-dimensional representations of $SU(2)$

$$\Psi[\vec{A}_U] = e^{i\varphi[\vec{A}_U]} \Psi[\vec{A}] \quad (\text{eigenstate of } U), \quad U_3 = U_2 U_1$$

$$e^{i\varphi[\vec{A}_{U_3}]} = e^{i\varphi[\vec{A}_{U_2}]} e^{i\varphi[\vec{A}_{U_1}]} \implies e^{i\varphi[\vec{A}_{U_2 U_1}]} - e^{i\varphi[\vec{A}_{U_1 U_2}]} = 0$$

$\implies \Psi'[\vec{A}]$ is gauge invariant

Throw states not satisfying $(*)$ out of the Hilbert space
→ CP conserved

Gauß' law

For $\Omega(\vec{x})$ an infinitesimal generator of gauge transformations

→ Noether charge:

$$\begin{aligned} Q(\Omega) &= \frac{1}{g} \int d^3x \operatorname{tr} \left[\Pi^i (D^i \Omega) \right] = \int_V d^3x \operatorname{tr} \left[\left(-E^i + \frac{g^2}{8\pi^2} \theta B^i \right) D^i \Omega \right] \\ &= \int d^3x \operatorname{tr} \left[\Omega D^i \left(E^i - \frac{g^2}{8\pi^2} \theta B^i \right) \right] + \int_{\partial V} da^i \operatorname{tr} \left[\Omega \left(-E^i + \frac{g^2}{8\pi^2} \theta B^i \right) \right] \end{aligned}$$

For $\Omega(\vec{x}) = 0$ when $\vec{x} \in \partial V$ and since Ψ' is gauge invariant

→ Gauß' law: $\vec{D} \cdot \vec{E} \Psi'[\vec{A}] = 0$

Usually, the argument is made the other way around: Impose Gauß' law to throw states out of the Hilbert space

Since $[Q(\Omega), W[\vec{A}]] = 0$ for $\Omega(\vec{x}) = 0$ when $\vec{x} \in \partial V$ this also holds when $\Psi'[\vec{A}] \rightarrow e^{i\tilde{\theta} W[\vec{A}]} \Psi'[\vec{A}]$, so imposing Gauß' law does not fix $\tilde{\theta}$, does not tell us about large gauge transformations

Nondiagonal basis

Redefining derivatives w.r.t. \vec{A} as

$$\vec{D}_{\vec{A}} \Psi[\vec{A}] = i \left(\frac{\delta}{i\delta\vec{A}} - (\theta - \theta_{\Pi}) \frac{g}{8\pi^2} \vec{B} \right) \Psi[\vec{A}]$$

corresponds to a canonical transformation of the momentum operator.

Induces translation as

$$T[\Delta\vec{A}] \Psi[\vec{A}] = e^{-i(\theta - \theta_{\Pi})(W[\vec{A} + \Delta\vec{A}] - W[\vec{A}])} \Psi[\vec{A} + \Delta\vec{A}]$$

For a shift $\Delta\vec{A}_{\text{gauge}}$ corresponding to a *general* gauge transformation: gauge invariant

$$T[\Delta\vec{A}_{\text{gauge}}] \Psi[\vec{A}] = \Psi[\vec{A}] \quad \text{if} \quad \Psi[\vec{A}] = e^{i(\theta - \theta_{\Pi})W[\vec{A}]} \Psi_{\text{g.i.}}[\vec{A}]$$

Agrees with reasoning & result in the diagonal basis

$\theta - \theta_{\Pi}$ in $\Psi_{\theta - \theta_{\Pi}}$ is pinned to $\theta - \theta_{\Pi}$ in H so that CP is conserved

Conclusion

There is no CP violation in QCD.

Challenges to the standard calculation and resolutions:

- Taking $T \rightarrow \infty$ after summing over sectors corresponds to an inequivalent deformation of the integration contour

Maintain contour and order of limits

- No point compactification/topology in temporal gauge (w/o extra constraint)

Identify the Hilbert space with gauge invariant observables for all gauge transformations on spatial hypersurfaces

- θ -vacua are not properly normalizable

Above Hilbert space allows to restrict inner product to integrate over each physical configuration one time and one time only

THANK YOU!