


Massive graviton dark matter searches with atom interferometers

John Carlton


john.carlton@kcl.ac.uk






arXiv: 1911.11755

Atom Interferometer Observatory and Network



Atom interferometry



Consider a 2-level atom



Consider a 2-level atom

Photon Absorption:


Mom. kick



Consider a 2-level atom



Photon Absorption:

Mom. kick





Stimulated Emission:


Mom. kick



Rabi oscillations



Interferometer sequence



Mach-Zehnder
interferometer




Image atom fringes
and measure phase

$$\phi_{\text{MZ}} = kgT^2$$

Leading order phase depends
on gravitational acceleration

What are we sensitive to?


What can we measure?

$$\phi_{\text{MZ}} = kgT^2$$

Atom-light
interactions


Gravitational field

Time




Atom gradiometer

Gradiometer phase
 $\Delta\phi = \phi_1 - \phi_2$




Atom cloud



Gradiometer phase


$$\Delta\phi = \phi_1 - \phi_2$$




Longer baseline L

Longer time of flight T

More sensitivity $\Delta\phi$




Gravitational waves



GW strain modifies laser propagation


$$h \sim \frac{\delta L}{L} \sim \frac{\delta T}{T}$$

Change in pulse timings affects phase




Gravitational waves

- ❖ ‘Mid-band’ sensitivity between LIGO and LISA.




Dark matter


A lot of parameter space!




A lot of parameter space!



A lot of parameter space!



A classical ULDM field




Ultralight mass means a high occupation number

Can describe as a classical field

$$\varphi(t, \mathbf{x}) \sim \cos(\omega_\varphi t - \mathbf{k}_\varphi \cdot \mathbf{x})$$

Frequency given by ULDM mass
(with small velocity correction)

$$\omega_\varphi \simeq m_\varphi \left(1 + \frac{v^2}{2} \right)$$




Atoms in a scalar ULDM field

$$\alpha(t, \mathbf{x}) \approx \alpha \left[1 + d_e \sqrt{4\pi G_N} \varphi(t, \mathbf{x}) \right],$$


$$m_e(t, \mathbf{x}) = m_e \left[1 + d_{m_e} \sqrt{4\pi G_N} \varphi(t, \mathbf{x}) \right]$$

$$\delta\phi \sim \delta\omega \sim \varphi(t, x)$$





Two sensitivity channels


Interrogation time (GWs)



Atomic transition frequency (Scalar ULDM)




What about spin-2?




Massive gravity field theory

Let's consider a massive spin-2 ultra-light field $\varphi_{\mu\nu}$



Massive gravity field theory

Let's consider a massive spin-2 ultra-light field $\varphi_{\mu\nu}$



Express as irreducible fields:


$$\varphi_{00} = \Psi$$

$$\varphi_{0i} = u_i + \partial_i v$$

$$\varphi_{ij} = \varphi_{ij}^{\text{TT}} + 2\partial_{(i} A_{j)} + \partial_i \partial_j \sigma + \delta_{ij} \pi$$

Massive gravity field theory

Let's consider a massive spin-2 ultra-light field $\varphi_{\mu\nu}$



Express as irreducible fields:

$$\varphi_{00} = \Psi$$

Tensor

$$\varphi_{0i} = u_i + \partial_i v$$

Vector

$$\varphi_{ij} = \varphi_{ij}^{\text{TT}} + 2\partial_{(i} A_{j)} + \partial_i \partial_j \sigma + \delta_{ij} \pi$$

Scalar

Three normalised fields

Tensor $\mathcal{L}_t = \frac{1}{2} (\tilde{\varphi}_{ij} \square \tilde{\varphi}_{ij} - m_t^2 \tilde{\varphi}_{ij} \tilde{\varphi}_{ij})$

Vector $\mathcal{L}_v = \frac{1}{2} (\tilde{A}_i \square \tilde{A}_i - m_v^2 \tilde{A}_i \tilde{A}_i)$

Scalar $\mathcal{L}_s = \frac{1}{2} (\tilde{\pi} \square \tilde{\pi} - m_s^2 \tilde{\pi}^2)$

Coupling to light and matter

$$\mathcal{L}_{\text{int}} = \kappa^\phi \varphi^{\mu\nu} \mathcal{O}_{\mu\nu}$$



Symmetric Standard Model operator

Coupling to light and matter

$$\mathcal{L}_{\text{int}} = \kappa^\phi \varphi^{\mu\nu} \mathcal{O}_{\mu\nu} \rightarrow \begin{array}{ccc} \text{Tensor} & \text{Vector} & \text{Scalar} \\ \kappa_t \varphi^{ij} \mathcal{O}_{ij}^t + \kappa_v \varphi^{0i} \mathcal{O}_{0i}^v + \kappa_s \varphi^{00} \mathcal{O}^s & & \end{array}$$

Coupling to light and matter

$$\mathcal{L}_{\text{int}} = \kappa^\phi \varphi^{\mu\nu} \mathcal{O}_{\mu\nu} \rightarrow \kappa_t \varphi^{ij} \mathcal{O}_{ij}^t + \kappa_v \varphi^{0i} \mathcal{O}_{0i}^v + \kappa_s \varphi^{00} \mathcal{O}^s$$

Tensor Vector Scalar

↓ ↓

Non-relativistic limit

$$\frac{\alpha}{M_{\text{Pl}}} \varphi_{ij}^{\text{TT}} F^{i\sigma} F_\sigma^j \quad \frac{\beta}{M_{\text{Pl}}} \tilde{\pi} (F^2 + m_\psi \bar{\psi} \psi)$$

Coupling to light and matter

$$\mathcal{L}_{\text{int}} = \kappa^\phi \varphi^{\mu\nu} \mathcal{O}_{\mu\nu} \rightarrow \kappa_t \varphi^{ij} \mathcal{O}_{ij}^t + \kappa_v \varphi^{0i} \mathcal{O}_{0i}^v + \kappa_s \varphi^{00} \mathcal{O}^s$$

Tensor Vector Scalar

\downarrow Non-relativistic limit \downarrow


$$\frac{\alpha}{M_{\text{Pl}}} \varphi_{ij}^{\text{TT}} F^{i\sigma} F_\sigma^j \quad \frac{\beta}{M_{\text{Pl}}} \tilde{\pi} (F^2 + m_\psi \bar{\psi} \psi)$$

In all theories Only in Lorentz violating theories!

Coupling to light and matter

Tensor modes

$$\frac{\alpha}{M_{\text{Pl}}} \varphi_{ij}^{\text{TT}} F^{i\sigma} F_\sigma^j$$




Coupling to light and matter

Tensor modes

$$\frac{\alpha}{M_{\text{Pl}}} \varphi_{ij}^{\text{TT}} F^{i\sigma} F_\sigma^j$$


Bends space-time
like gravitational
waves!



Scalar modes

$$\frac{\beta}{M_{\text{Pl}}} \tilde{\pi}(F^2 + m_\psi \bar{\psi}\psi)$$

Modifies atomic
energy levels just
like scalar ULDM!




What can we measure?

$$\phi_{\text{MZ}} = kgT^2$$


Atom-light
interactions

Gravitational field

Time




What can we measure?



Projected detection limits

10m, 100m and 1km example atom interferometers

Isotope	L [m]	T [s]	n	Δr [m]	S_n [Hz^{-1}]
^{87}Sr	10	0.74	1000	5	10^{-8}
^{87}Sr	100	1.4	1000	90	10^{-10}
^{87}Sr	1000	1.4	1000	980	0.09×10^{-10}




Projected detection limits

10m, 100m and 1km example atom interferometers

Assume Lorentz invariance so only **tensor** modes contribute

$$\varphi_{\mu}^{\mu} = 0$$




Projected detection limits

Leading constraints on tensor mode come from ‘fifth force’ experiments

$$\delta V_{\text{Newt}} \propto \alpha^2 e^{-m_{\tilde{\varphi}} r}$$

In this range, from lunar laser ranging



Projected detection limits

Leading constraints on tensor mode come from ‘fifth force’ experiments


$$\delta V_{\text{Newt}} \propto \alpha^2 e^{-m_{\tilde{\varphi}} r}$$

In this range, from lunar laser ranging

Higuchi bound sets a lower bound for mass of spin-2 field

$$m^2 \geq 2H^2$$

Least stringent bound from BBN




Projected detection limits

Consider the tensor and scalar couplings independently in the Lorentz violating case.

$$m_t \neq m_s$$


$$\alpha \neq \beta$$

Scalar mode constrained by tests of equivalence principle



Advantages of networking!

AION plans to network with MAGIS-100 to enhance sensitivity in ULDM/GW searches.




Advantages of networking!


AION plans to network with MAGIS-100 to enhance sensitivity in ULDM/GW searches.

$$\varphi_{ij}^{\text{TT}}(t, \mathbf{x}) = \sum_{\lambda} \varphi_{0,\lambda}^{\text{TT}} e_{ij}^{\lambda}(\mathbf{k}_t) \cos(\omega_t t - \mathbf{k}_t \cdot \mathbf{x})$$

Distinguish dark matter models through directional dependence.



Progress towards a global network!



Summary

AION is an upcoming atom interferometer experiment, using quantum sensors for detecting ultralight dark matter and gravitational waves – in the ‘mid-band’ between LISA and LIGO.

Spin-2 ULDM can be probed by gravitational wave detectors – however, atom interferometers can detect it through two different channels without altering any of the experimental design!

A global network of atom interferometers will enhance these searches further, probing the directional dependence of the field and other couplings.

Backup


LMT pulses

Additional pulses
enhance sensitivity

$$n = 2$$


Atom gradiometer

$$\Delta\phi = \phi_1 - \phi_2$$




Scalar ULDM sensitivity

- ❖ ULDM mass/frequency sensitivity depends on T .



Phase shifts

$$\phi = \boxed{\phi_{\text{prop}}} + \boxed{\phi_{\text{sep}}} + \boxed{\phi_{\text{laser}}} = kgT^2$$




Spin-1 dark matter

B-L coupling, which generates
a ‘dark’ electric field

$$\Delta F_{B-L} \sim g_{B-L} \left(\frac{Z_1}{A_1} - \frac{Z_2}{A_2} \right) E_{B-L}$$

Probe with a dual-species interferometer



AION-10 sensitivity projections

$$d_{m_e}^{\text{best}} \sim \left(\frac{1}{T}\right)^{5/4} \frac{1}{C n \Delta r} \left(\frac{\Delta t}{N_a}\right)^{1/2} \left(\frac{1}{T_{\text{int}}}\right)^{1/4}$$

Handles to optimise (in order of priority):

$T \sim 1\text{s}$ (interrogation time)

$C \sim 0.1 - 1$ (contrast)


$n \sim 1000$ (LMT)

$\Delta r \sim \text{Al separation}$

$\Delta t \sim \text{sampling time}$

$N_a \sim \text{atoms in cloud}$

$T_{\text{int}} \sim 10^7\text{s}$ (integration time)




Massive gravity field theory

Let's consider a massive spin-2 ultra-light field $\varphi_{\mu\nu}$

Fierz-Pauli Lagrangian


$$\mathcal{L}_{\text{FP}} = \mathcal{L}_{\text{EH}} - \frac{1}{4}m^2 (\varphi_{\mu\nu}\varphi^{\mu\nu} - \varphi^2)$$

Lorentz invariant massive spin-2 field



Massive gravity field theory

Let's consider a massive spin-2 ultra-light field $\varphi_{\mu\nu}$



$$\mathcal{L}_{\text{FP}} = \mathcal{L}_{\text{EH}} - \frac{1}{4} \left(m_0^2 \varphi_{00}^2 + 2m_1^2 \varphi_{0i}^2 - m_2^2 \varphi_{ij}^2 + m_3^2 \varphi_i^i \varphi_j^j - 2m_4^2 \varphi_{00} \varphi_i^i \right)$$

Lorentz violating massive spin-2 field