



MAPPING NEUTRON STARS INSIDE AND OUT



PROF. ANNA WATTS
(UNIVERSITY OF AMSTERDAM)

THE NEUTRON STAR INTERIOR



FROM NUCLEAR PHYSICS TO TELESCOPE (AND BACK)






Figure: Adapted from Ray et al. 2019



NICER



Image: NASA

THE PULSE PROFILE MODELING TECHNIQUE



Physics and priors

- Hotspot properties (magnetic poles)
- Atmospheric beaming
- Relativistic ray-tracing in rotating NS space-time (mass, radius, spin)
- Distance and observer inclination
- Interstellar absorption
- Instrument response including calibration uncertainty
- Background
- May have priors on mass, distance, inclination from radio timing
- 'EOS-agnostic' priors on radius




Bayesian inference of model parameters (statistical sampling)

Mass and radius \Rightarrow dense matter
Hotspot properties \Rightarrow magnetic pole/field geometry

PULSE PROFILE DATA SETS

PSR J0030+0451


- Bright source
- Isolated system



Riley et al. 2019, Miller et al. 2019
Salmi et al. 2023, Vinciguerra et al. 2023,
2024

PSR J0740+6620


- Faint source
- Binary - mass $2.1M_{\odot}$



Riley et al. 2021, Miller et al. 2021
Salmi et al. 2022, 2023, 2024a,
Dittmann et al. 2024, Hoogkamer et al. 2025

PSR J0437-4715


- Closest and brightest source
- Binary: mass $1.4M_{\odot}$




Choudhury et al. 2024

MAPPING THE HOT MAGNETIC POLES


PSR J0030+0451



PSR J0740+6620



PSR J0437-4715




As seen from Earth

Results from the X-PSI pipeline (<https://github.com/xpsi-group/xpsi>, also Riley et al. 2023 JOSS)


Maps from Riley et al. 2019, Riley et al. 2021, Choudhury et al. 2024 + Morsink/NASA

NON-DIPOLAR MAGNETIC FIELD?



Credit: NASA's Goddard Space Flight Center/Harding, Kalapotharakos, Wadiasingh.

MASSSES AND RADII



PSR J0437-4715


- Very well-constrained mass (Reardon et al. 2024)
- AGN in field of view \Rightarrow complex background
- PPM analyses to date
 - 2017-2021 data set (Choudhury et al. 2024)
- Consistent with softer EOS, and with GW170817

- Hoogkamer et al. 2026 (PRD in press, arXiv:2510.27619) studied this data set with EOS-informed priors. Found new narrow geometric mode. Higher likelihood, but physically less likely since inconsistent with multi-wavelength pulsar data/models.

- In progress: analysis of extended data set (Choudhury++, Stammler++)

- Miller et al 2025, arXiv:2512.08790: analysis including additional model component

MASSES AND RADII




PSR J0437-4715

PSR J0740+6620

- Well-constrained, high mass (Cromartie et al. 2020, Fonseca et al. 2021)
- PPM analyses to date:
 - 2018-2020 data set (Riley et al. 2021, Miller et al. 2021)
 - 2018-2021 data set, cross-testing background treatments (Salmi et al. 2022)
 - 2018-2022 data set (Salmi et al. 2024a, Dittmann et al. 2024)
- Very softest EOS disfavored
- Hoogkamer et al. 2025 (PRD) tested sampler dependence/confirmed convergence of results from Salmi et al. 2024a
- Hoogkamer et al. 2026 (PRD in press) tested same data set with EOS-informed priors.

MASSES AND RADII




PSR J0437-4715

PSR J0740+6620

PSR J0030+0451

- No mass prior (not in a binary)
- PPM analyses to date:
 - 2017-2018 data set, no background constraints (Riley et al. 2019, *Miller et al. 2019*)
 - Replication of X-PSI analysis by Afle et al. (2023)
 - 2017-2018 data set, with background constraints and improved sampler settings (Vinciguerra et al. 2024)
- Shown here is solution from V24 without background constraints.

MASSES AND RADII




PSR J0437-4715

PSR J0740+6620

PSR J0030+0451

- No mass prior (not in a binary)
- PPM analyses to date:
 - 2017-2018 data set, no background constraints (Riley et al. 2019, *Miller et al. 2019*)
 - Replication of X-PSI analysis by Afle et al. (2023)
 - 2017-2018 data set, with background constraints and improved sampler settings (Vinciguerra et al. 2024)
- With background, there are multimodal solutions! Some consistent with PSR J0437-4715 (Model 1).

MASSES AND RADII




PSR J0437-4715

PSR J0740+6620

PSR J0030+0451

- No mass prior (not in a binary)
- PPM analyses to date:
 - 2017-2018 data set, no background constraints (Riley et al. 2019, *Miller et al. 2019*)
 - Replication of X-PSI analysis by Afle et al. (2023)
 - 2017-2018 data set, with background constraints and improved sampler settings (Vinciguerra et al. 2024)
- With background, there are multimodal solutions! Some consistent with PSR J0437-4715 (Model 1).
- Analysis of full data set with background constraints now complete (Kini et al. in prep).

MASSES AND RADII



SR J0437-4715

SR J0740+6620

SR J0030+0451

No mass prior (not in a binary)


PPM analyses to date:

- 2017-2018 data set, no background constraints (Riley et al. 2019, *Miller et al. 2019*)
- Replication of X-PSI analysis by Afle et al. (2023)
- 2017-2018 data set, with background constraints and improved sampler settings (Vinciguerra et al. 2024)

With background, there are multimodal solutions!
Some consistent with PSR J0437-4715 (Model 1).

Analysis of full data set with background constraints now complete (Kini et al. in prep).

MASSES AND RADII




PSR J0437-4715

PSR J0740+6620

PSR J0030+0451

- No mass prior (not in a binary)
- PPM analyses to date:
 - 2017-2018 data set, no background constraints (Riley et al. 2019, *Miller et al. 2019*)
 - Replication of X-PSI analysis by Afle et al. (2023)
 - 2017-2018 data set, with background constraints and improved sampler settings (Vinciguerra et al. 2024)
- With background, there are multimodal solutions! Some consistent with PSR J0437-4715 (Model 1).
- Analysis of full data set with background constraints now complete (Kini et al. in prep).

MASSES AND RADII




PSR J0437-4715

PSR J0740+6620

PSR J0030+0451

- No mass prior (not in a binary)
- PPM analyses to date:
 - 2017-2018 data set, no background constraints (Riley et al. 2019, *Miller et al. 2019*)
 - Replication of X-PSI analysis by Afle et al. (2023)
 - 2017-2018 data set, with background constraints and improved sampler settings (Vinciguerra et al. 2024)
- With background, there are multimodal solutions! Some consistent with PSR J0437-4715 (Model 1).
- Analysis of full data set with background constraints now complete (Kini et al. in prep).

MASSES AND RADII




PSR J0437-4715

PSR J0740+6620

PSR J0030+0451

- No mass prior (not in a binary)
- PPM analyses to date:
 - 2017-2018 data set, no background constraints (Riley et al. 2019, *Miller et al. 2019*)
 - Replication of X-PSI analysis by Afle et al. (2023)
 - 2017-2018 data set, with background constraints and improved sampler settings (Vinciguerra et al. 2024)
- With background, there are multimodal solutions! Some consistent with PSR J0437-4715 (Model 1).
- Analysis of full data set with background constraints now complete (Kini et al. in prep).

MASSES AND RADII




PSR J0437-4715

PSR J0740+6620


PSR J0030+0451

PSR J1231-1411

- Unusual pulse profile (very weak 2nd pulse)



MASSES AND RADII



PSR J0437-4715


PSR J0740+6620

PSR J0030+0451


PSR J1231-1411

- Unusual pulse profile (very weak 2nd pulse)
- Weakly-constrained mass prior
- PPM analysis to date:
 - 2017-2023 data set (Salmi et al. 2024b)
- Restricted radius prior required for convergence (so be careful using this in EOS analyses!)
- We can find good solutions that are consistent with EOS inferred from the other sources
- Work underway to determine why this source is so challenging (see also Qi et al. 2025)

MASSES AND RADII



MASSES AND RADII



PSR J0437-4715

PSR J0740+6620


PSR J0030+0451

PSR J1231-1411

PSR J0614-3329

- Constrained mass prior from MeerKAT PTA (Miles et al. 2025)
- PPM analysis to date
 - 2017 – 2023 data set (Mauviard et al. 2025)
- Geometry – 1 polar, 1 equatorial hotspot
- All (geometric) models favor median radii in 9 - 11km range

CONSISTENCY WITH GW?



PSR J0437-4715

PSR J0740+6620


PSR J0030+0451

PSR J1231-1411

PSR J0614-3329


GW170817: LSC and VSC 2018

EQUATION OF STATE INFERENCE



- Start with our inferred mass-radius posteriors
- Select an EOS model (with parameters and priors on those parameters)
- Infer EOS model parameters and central densities -> Inferred EOS
- This then translates into an inferred mass-radius **relation**


EOS INFERENCE (EXAMPLE)



EOS model

- BPS outer crust
- cEFT up to 1.5 saturation density (Keller et al. 2023)
- Piecewise polytrope extensions at high density
- Full description and alternative models in Rutherford et al. 2024, see also Mauviard et al. 2025.
- Prior space for this model excludes high radii, favours radii around 12-13km.

EOS INFERENCE (EXAMPLE)



EOS model

- BPS outer crust
- cEFT up to 1.5 saturation density (Keller et al. 2023)
- Piecewise polytrope extensions at high density
- Full description and alternative models in Rutherford et al. 2024, see also Mauviard et al. 2025.
- Prior space for this model excludes high radii, favours radii around 12-13km.
- Posteriors once NICER and GW data are included (Rutherford et al. 2024, see Mauviard et al. 2025 for an update that includes J0614).

NICER REPAIR – EVA 91, 16 JAN 2025




Image: NASA

June 2025: NICER
operations suspended



Image: NASA

PPM FOR ACCRETING/BURSTING NEUTRON STARS




- Spots are complex (Das et al. 24, 25).
- Understanding the disk is important (Dorsman et al. 25, 26)
- Polarimetry can help! (Salmi et al. 2025)
- Joint NICER/IXPE analysis for SGRA J1444 underway (Dorsman, Salmi + in prep)




Variability a major challenge, but solvable (Kini et al. 23,24a)
First results find small R (Kini et al. 24b) – but model needs work.

AMPs a major target for eXTP
(launch 2030)

Improved surface pattern models for all source types



EOS analysis: follow on (e.g. Rutherford et al. 2024, Mauviard et al. 2025) and now more integrated in PPM (Hoogkamer et al. 2026)



NICER legacy - 8 yrs of amazing data!



New X-ray telescopes: eXTP (2030), NewAthena



Better priors from radio pulsar timing

SUMMARY

- Results to date for rotation-powered millisecond pulsars point to a soft to medium stiffness EOS, and non-dipolar fields.
- More data and sources to come.
- PPM pipelines are evolving e.g more physical surface patterns, EOS informed priors.....
- PPM is now being applied to accreting neutron stars, and we are preparing for new telescopes (eXTP, NewAthena)

