

Neutrino propagation in the presence of nonstandard interactions in binary neutron star mergers


Amélie Chatelain

Laboratoire AstroParticule et Cosmologie, Paris Diderot
A. Chatelain, C. Volpe, Arxiv:17xx:xxxx

October 13, 2017


- ① Framework
- ② Impact of nonstandard interactions in BNS mergers
- ③ Conclusions and perspectives

Binary Neutron Star mergers



[Perego et al., Mon.Not.Roy.Astron.Soc.

443, 2014]



→ How can nonstandard interactions impact neutrino flavor conversions in this environment ?

Neutrino flavor conversions in BNS : the model

- Two neutrino flavors in the mean-field approximation.
- Two evolution equations : $i\dot{\rho} = [H, \rho]$ and $i\dot{\bar{\rho}} = [\bar{H}, \bar{\rho}]$.

$$H = H_{\text{vac}} + H_{\text{mat}} + H_{\text{self}}$$

$H_{\text{vac}} = \omega \begin{pmatrix} -c_{2\theta} & s_{2\theta} \\ s_{2\theta} & c_{2\theta} \end{pmatrix}$

$H_{\text{mat}} = \lambda Y_e \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$


Astrophysical simulations

The Matter Neutrino Resonance

- $L_{\bar{\nu}_e} > L_{\nu_e}$: possible MSW-like cancellation between the matter and ν self-interaction term → **Matter Neutrino resonance**.

$$H_{ee} - H_{xx} = \lambda Y_e + H_{\text{self},ee} - H_{\text{self},xx} - 2\omega c_{2\theta} \approx 0$$

- Adiabaticity increased by nonlinear feedback.



[Malkus, Kneller, McLaughlin, Surman, PRD86,2012] [Malkus, Friedland, McLaughlin, 1403.5797] [Malkus, McLaughlin, Surman, PRD93,2016] [Vaananen, McLaughlin, PRD93, 2016] [Zhu, Perego, McLaughlin, PRD94, 2016] [Frensel, Wu, Volpe, Perego, PRD95, 2017]

Nonstandard interactions

- Matter-Neutrino NSI. Possible origin of anomalies, searched in various experiments (oscillation, scattering, ...).
- Constraints :

$$\begin{pmatrix} |\epsilon_{ee}| < 2.5 & |\epsilon_{e\mu}| < 0.21 & |\epsilon_{e\tau}| < 1.7 \\ & |\epsilon_{\mu\mu}| < 0.05 & |\epsilon_{\mu\tau}| < 0.2 \\ & & |\epsilon_{\tau\tau}| < 9.0 \end{pmatrix}$$

- New term to our Hamiltonian.

$$H_{\text{NSI}} = \lambda \begin{pmatrix} \epsilon_{ee} & \epsilon_{e\tau} \\ \epsilon_{\tau e} & \epsilon_{\tau\tau} \end{pmatrix} \rightarrow \lambda \begin{pmatrix} \frac{Y_\odot - Y_e}{Y_\odot} \delta \epsilon^n & (3 + Y_e) \epsilon_0 \\ (3 + Y_e) \epsilon_0^* & 0 \end{pmatrix}$$

$$|\delta \epsilon^n| \leq \mathcal{O}(1 - 10)$$

$$|\epsilon_0| \leq \mathcal{O}(0.1 - 1)$$

- Explored in Sun and SNe. [Friedland, Lunardini, Pena-Garay, PLB594, 2004]
 [Esteban-Pretel, Tomas, Valle, PR76, 2007] [Blennow, Mirizzi, Serpico, PRD78, 2008]
 [Stapleford, Vaananen, Kneller, McLaughlin, Shapiro, PRD94, 2016] [Sen, Dighe, 2017]


→ What are the effects of NSI in binary neutron star mergers ?

Matter Neutrino Resonance modified


$$\underbrace{\lambda \left[Y_e + \frac{Y_{\odot} - Y_e}{Y_{\odot}} \delta \epsilon^n \right]}_{V_M} + H_{\text{self},ee} - H_{\text{self},xx} - 2\omega c_{2\theta} \approx 0$$

Can modify the **location** and **adiabaticity** of the MNR.

MNR
without
NSI






MNR
with
NSI



The I-resonance

$$\underbrace{\lambda \left[Y_e + \frac{Y_\odot - Y_e}{Y_\odot} \delta \epsilon^n \right]}_{V_M} \approx 2\omega c_{2\theta} - (H_{\text{self},ee} - H_{\text{self},xx}) \approx 0$$

- Self-interaction : no effect on location or adiabaticity of the resonance. [Esteban-Pretel, Tomas, Valle, PR76, 2007]
- Can be localized extremely close to the neutrinosphere.
- Creates similar transformations for neutrinos and antineutrinos.



I-resonance in the presence of strong self-interaction

- When the self-interaction potential is very strong ($\mu > \lambda$), the I-resonance can still occur as a synchronized MSW resonance.
- Define the synchronized mode

$$\vec{J} = \sum_{\alpha=e,x} \int_0^\infty dp \left(G_{\nu_\alpha} j_{\nu_\alpha}(p) \vec{P}_{\nu_\alpha}(p) - G_{\bar{\nu}_\alpha} j_{\bar{\nu}_\alpha}(p) \vec{\bar{P}}_{\bar{\nu}_\alpha}(p) \right)$$

$$\partial_r \vec{J} \approx \left(\omega_{\text{sync}} \vec{B}_0 + \vec{B}_{\text{mat}} \right) \times \vec{J} + \frac{\partial_r \mu}{\sqrt{2} G_F} \hat{J}$$

- MSW-like resonance condition for this mode :





$$\omega_{\text{sync}} c_{2\theta} = V_M$$

$$\omega_{\text{sync}} = \frac{\sqrt{2} G_F \Delta m^2 F_1(0) F_3(0)}{2\mu F_2^2(0)} \left[\frac{L_{\nu_e} G_{\nu_e}}{R_{\nu_e}^2 \langle E_{\nu_e} \rangle^2} + \frac{L_{\bar{\nu}_e} G_{\bar{\nu}_e}}{R_{\bar{\nu}_e}^2 \langle E_{\bar{\nu}_e} \rangle^2} - 2 \frac{L_{\nu_x} G_{\nu_x}}{R_{\nu_x}^2 \langle E_{\nu_x} \rangle^2} \right] \propto \frac{1}{\mu}$$

I-resonance as synchronized-MSW : proof of principle

$$\omega_{\text{sync}} c_{2\theta} = V_M$$




I-resonance : can be found in the presence of a strong self-interaction potential, as a **synchronized** MSW resonance.




Combination MNR and I-resonance

Presence of I-resonance : modify the flavor content.

- Can trigger the Matter Neutrino Resonance.
- Can encounter multiple resonances, with different patterns.




Combination MNR and I-resonance



Impact on nucleosynthesis : evolution of Y_e

$$V_M = \lambda \left[Y_e + \frac{Y_\odot - Y_e}{Y_\odot} \delta \epsilon^n \right] \approx 0$$

Plot : I resonance condition for $\delta \epsilon^n \in [-0.9, -0.2]$.



I-resonances can be found for a large range of NSI parameters.

Conclusions and perspectives

- Neutrino flavor conversions in BNS mergers : lots of on-going investigations.
- **Nonstandard Interactions** : have a significant impact on flavor conversions.
 - Can shift the usual Matter Neutrino Resonance.
 - Can create an **Inner resonance** very close to the neutrinosphere, which can also be synchronized.
 - Can create combination and different patterns of the two resonances.
 - Impact on nucleosynthesis : under study ! "*Neutrino propagation in the presence of nonstandard interactions in binary neutron star mergers*", Chatelain, Volpe, Arxiv:17xx:xxxx

Conclusions and perspectives

- Neutrino flavor conversions in BNS mergers : lots of on-going investigations.
- **Nonstandard Interactions** : have a significant impact on flavor conversions.
 - Can shift the usual Matter Neutrino Resonance.
 - Can create an **Inner resonance** very close to the neutrinosphere, which can also be synchronized.
 - Can create combination and different patterns of the two resonances.
 - Impact on nucleosynthesis : under study ! "*Neutrino propagation in the presence of nonstandard interactions in binary neutron star mergers*", Chatelain, Volpe, Arxiv:17xx:xxxx

Thank you !